

THE NON-SELFADJOINT APPROACH TO THE HAO-NG ISOMORPHISM

ELIAS G. KATSOULIS AND CHRISTOPHER RAMSEY

ABSTRACT. In an earlier work, the authors proposed a non-self-adjoint approach to the Hao-Ng isomorphism problem for the full crossed product, depending on the validity of two conjectures stated in the broader context of crossed products for operator algebras. By work of Harris and Kim, we now know that these conjectures in the generality stated may not always be valid. In this paper we show that in the context of hyperrigid tensor algebras of C^* -correspondences, each one of these conjectures is equivalent to the Hao-Ng problem. This is accomplished by studying the representation theory of non-selfadjoint crossed products of C^* -correspondence dynamical systems; in particular we show that there is an appropriate dilation theory. A large class of tensor algebras of C^* -correspondences, including all regular ones, are shown to be hyperrigid. Using Hamana's injective envelope theory, we extend earlier results from the discrete group case to arbitrary locally compact groups; this includes a resolution of the Hao-Ng isomorphism for the reduced crossed product and all hyperrigid C^* -correspondences. A culmination of these results is the resolution of the Hao-Ng isomorphism problem for the full crossed product and all row-finite graph correspondences; this extends a recent result of Bedos, Kaliszewski, Quigg and Spielberg.

1. INTRODUCTION

Let $((X, \mathcal{C}), \mathcal{G}, \alpha)$ be a C^* -correspondence dynamical system where \mathcal{G} is a locally compact group and α is a generalized gauge action. This action can be extended uniquely to the Cuntz-Pimsner algebra \mathcal{O}_X . The *Hao-Ng isomorphism problem* asks whether

$$\mathcal{O}_X \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{O}_{X \rtimes_{\alpha} \mathcal{G}}$$

in the reduced or full crossed products. This problem is named after Hao and Ng who proved the validity of this formula when \mathcal{G} is amenable

2010 *Mathematics Subject Classification.* 46L07, 46L08, 46L55, 47B49, 47L40, 47L65.

Key words and phrases: C^* -correspondence, crossed product, Cuntz-Pimsner algebra, tensor algebra, Hao-Ng isomorphism, C^* -envelope, operator algebra.

[13, Theorem 2.10]. However, this formula was first studied by Abadie in the context of Takai duality for equivalence bimodules. Indeed, in Abadie's proof for the Takai duality, the Hao-Ng isomorphism forms the crucial step of the proof and corresponds to the key isomorphism of [32, Lemma 7.2] in the classical case. In general, the Hao-Ng isomorphism has proved to be a significant stimulant to research as versions of it appear in many different context, e.g. in Schafhauser's work [31] on AF-embedability, or in Deaconu's work [7, 8] on group actions on graph C^* -algebras. In its full generality, the problem remains open and under investigation by several authors [3, 16, 17, 22, 25].

The authors initiated a study in [20, 18, 21] of non-selfadjoint crossed products of operator algebra dynamical systems $(\mathcal{A}, \mathcal{G}, \alpha)$ where α acts by completely isometric isomorphisms of \mathcal{A} . The main thrust of [20, Chapter 7] and [18] is that the Hao-Ng isomorphism problem can and should be thought of as a non-selfadjoint problem. For the reduced crossed product this kind of approach has been and continues to be quite successful. For instance, we now know that the Hao-Ng isomorphism for the reduced crossed product holds for all discrete groups [18], a fact that resolves an open problem from [3] (and more is accomplished in this paper).

The Hao-Ng isomorphism for the *full* crossed product seems to be a much harder problem. In [20] we proposed the following line of attack. (See just above Theorem 7.9 in [20].) First one verifies

$$(1) \quad C_{\text{env}}^*(\mathcal{A} \rtimes_{\alpha} \mathcal{G}) \simeq C_{\text{env}}^*(A) \rtimes_{\alpha} \mathcal{G}$$

in the full crossed product case for an arbitrary non-selfadjoint dynamical system $(\mathcal{A}, \mathcal{G}, \alpha)$; this is Problem 1 in [20]. Subsequently, one solves Problem 2 in [20] by showing that all relative crossed products coincide.¹ Assuming that both problems have been resolved in the positive, now one specializes on tensor algebra dynamical systems and obtains

$$(2) \quad \mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha} \mathcal{G}}^+$$

by invoking the solution of Problem 2 and the remarks following [20, Theorem 7.13]. Recalling that $C_{\text{env}}^*(\mathcal{T}_X^+) = \mathcal{O}_X$, one recovers now the Hao-Ng isomorphism by combining equations (1) and (2). Note that even though a positive answer for *both* Problems 1 and 2 leads to a positive resolution for the Hao-Ng isomorphism, the exact relation of each one of these problems with the Hao-Ng isomorphism was never clarified in [20].

¹This also establishes that (1) is valid for the relative crossed product $\mathcal{A} \rtimes_{C_{\text{env}}^*(\mathcal{A}), \alpha} \mathcal{G}$, as it coincides with $\mathcal{A} \rtimes_{\alpha} \mathcal{G}$.

The central result of this paper, Theorem 4.9, clarifies that relation and shows that the Hao-Ng problem actually leads to equivalent statements in non-selfadjoint operator algebra theory, whose validity or refutation will therefore resolve the isomorphism. Specifically, for a large class of C^* -correspondences, including all regular ones, the validity of (1) for $\mathcal{A} = \mathcal{T}_X^+$ is equivalent to the validity of the Hao-Ng isomorphism $\mathcal{O}_X \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{O}_{X \rtimes_{\alpha} \mathcal{G}}$. In addition, we show that the validity of the Hao-Ng isomorphism $\mathcal{O}_X \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{O}_{X \rtimes_{\alpha} \mathcal{G}}$ is equivalent to the fact that all relative crossed products for $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ coincide, where α is a generalized gauge action. (For a general C^* -correspondence X we show that the above statements are successively weaker.)

Theorem 4.9 relates to exciting new work by Harris and Kim [14]. Indeed these authors have answered both Problems 1 and 2 from [20, Chapter 7] by producing finite dimensional, hyperrigid dynamical systems $(\mathcal{A}, \mathcal{G}, \alpha)$ with distinct relative crossed products and failing (1). However the examples of Harris and Kim [14] do not concern tensor algebras of C^* -correspondences and so the Hao-Ng problem remains open. Theorem 4.9 shows now that the resolution of the Hao-Ng problem will lead to or will follow from the existence or the absence of Harris-Kim type examples but in the realm of tensor algebras. Needless to say that the quest for such examples, or the refutation of their existence, becomes now a project of high priority.

To test our new results, we study the Hao-Ng isomorphism for a class of C^* -correspondences that plays a central role in the theory: graph C^* -correspondences. In Theorem 5.4 we show that the Hao-Ng isomorphism problem is true in the case of row-finite graph correspondences, thus showing that the crossed product of such a Cuntz-Krieger algebra is the Cuntz-Pimsner algebra of a crossed product (of a graph) correspondence. This is done by showing that in the case of a dynamical system $(\mathcal{A}, \mathcal{G}, \alpha)$ where \mathcal{A} is the tensor algebra of *any* graph, \mathcal{G} any locally compact group and α a generalized gauge action, all relative crossed products coincide. Then, Theorem 4.9 finishes the proof for row-finite graphs. Note that in the special case where \mathcal{G} is discrete, Theorem 5.4 has also been obtained independently by Bedos, Kaliszewski, Quigg and Spielberg using different methods [4, Corollary 6.8 and Remark 6.10]. It is worth mentioning here that Theorem 5.4 is essentially obtained by dilating representations in a wholly constructive manner and should prove of much interest to those who study the representation theory of C^* -correspondences. At the moment, the lack of a constructive dilation proof of $C_{\text{env}}^*(\mathcal{T}_X^+) = \mathcal{O}_X$ seems to be a barrier to establishing (2) for the full crossed product in general.

On the way to proving the above theorems we obtain several results of independent interest. First we resolve Problem 3 from our monograph [20, Chapter 8]. Specifically, we show if (X, \mathcal{C}) is a non-degenerate C^* -correspondence and $\alpha : \mathcal{G} \rightarrow (X, \mathcal{C})$ is the generalized gauge action of a locally compact group, then $\mathcal{T}_X^+ \rtimes_\alpha \mathcal{G}$ is necessarily the tensor algebra of some C^* -correspondence.

Another result of independent interest is Theorem 3.1, which identifies a large class of hyperrigid C^* -correspondences, i.e., C^* -correspondences whose tensor algebras are hyperrigid. Indeed our central Theorem 4.9 actually applies to all hyperrigid C^* -correspondences. To make that result usable, we show that any C^* -correspondence $(X, \mathcal{C}, \varphi_X)$ with $\varphi_X(\mathcal{J}_X)X = X$ is hyperrigid (here \mathcal{J}_X denotes Katsura's ideal). This includes all previous known examples of hyperrigid C^* -correspondences and many more, e.g., all regular ones.

An interesting byproduct of our techniques on the full crossed product version of the Hao-Ng problem, is the resolution of the same problem for the reduced crossed product and all hyperrigid C^* -correspondences. In [18] the first named author verified the Hao-Ng isomorphism for the reduced crossed product and all discrete groups. Because here we are addressing locally compact groups which may not be discrete, we have to use an approach different from that of [18]. In particular, the algebra \mathcal{A} does not embed in either $\mathcal{A} \rtimes_\alpha \mathcal{G}$ or $\mathcal{A} \rtimes_\alpha^r \mathcal{G}$ and so restricting a maximal map of the crossed product on the core algebra \mathcal{A} (as we did in [18]) is no longer an option. Instead we use Hamana's injective envelope theory, an approach towards the Hao-Ng isomorphism which is used in this paper for the first time. This approach was adopted after illuminating discussions with S. Echterhoff and we are grateful for that.

2. CROSSED PRODUCTS AND C^* -COVERS

Let $(\mathcal{A}, \mathcal{G}, \alpha)$ be an operator algebra dynamical system, meaning that \mathcal{A} is an approximately unital operator algebra and \mathcal{G} is a locally compact (Hausdorff) group acting continuously on \mathcal{A} by completely isometric automorphisms, $\alpha : \mathcal{G} \rightarrow \text{Aut}(\mathcal{A})$. The aim of this section is to better understand the relationship of α -admissible C^* -covers. Recall that a C^* -cover (\mathcal{C}, ι) of \mathcal{A} is a C^* -algebra \mathcal{C} and a complete isometry $\iota : \mathcal{A} \rightarrow \mathcal{C}$ such that $C^*(\iota(\mathcal{A})) = \mathcal{C}$.

The two nicest C^* -covers of \mathcal{A} are the “biggest” and the “smallest” covers $C_{\max}^*(\mathcal{A})$ and $C_{\text{env}}^*(\mathcal{A})$. These are defined by their universal properties. Namely, whenever (\mathcal{C}, ι) is a C^* -cover there are (unique)

surjective $*$ -homomorphisms $\varphi : C_{\max}^*(\mathcal{A}) \rightarrow \mathcal{C}$ and $\psi : \mathcal{C} \rightarrow C_{\text{env}}^*(\mathcal{A})$ such that $\varphi(a) = \iota(a)$ and $\psi(\iota(a)) = a$, for all $a \in \mathcal{A}$.

From [20], a C^* -cover (\mathcal{C}, ι) is called α -admissible if there exists a group representation $\beta : \mathcal{G} \rightarrow \text{Aut}(\mathcal{C})$ acting on \mathcal{C} by $*$ -automorphisms such that

$$\beta_g(\iota(a)) = \iota(\alpha_g(a)), \quad \forall g \in \mathcal{G}, a \in \mathcal{A}.$$

In [20, Lemma 3.3] we established that both $C_{\text{env}}^*(\mathcal{A})$ and $C_{\max}^*(\mathcal{A})$ are always α -admissible. However, in [20] we did not provide any examples of C^* -covers which fail to be α -admissible. We thank David Sherman for bringing this to our attention and asking us whether such covers do exist.

Proposition 2.1. *Not all C^* -covers are α -admissible.*

Proof. Let $\mathcal{C} = C(\mathbb{T}) \oplus M_2$ and $\iota : A(\mathbb{D}) \rightarrow \mathcal{C}$ be given by $z \mapsto z \oplus \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$. By von Neumann's inequality it is straightforward that ι is a complete isometry. Now

$$\begin{aligned} \iota(z) - \iota(z^2)\iota(z)^* &= (z \oplus \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}) - (z^2 \oplus \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix})(\bar{z} \oplus \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}) \\ &= 0 \oplus \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}. \end{aligned}$$

Thus, $C^*(\iota(A(\mathbb{D}))) = \mathcal{C}$ and (\mathcal{C}, ι) is a C^* -cover of $A(\mathbb{D})$.

Consider the Möbius transformation $\varphi(z) = \frac{z - \frac{1}{2}}{1 - \frac{z}{2}}$ which gives $\varphi \in \text{Aut}(\mathbb{D})$. From this define the dynamical system $(A(\mathbb{D}), \alpha, \mathbb{Z})$ where $\alpha_n(f) = f \circ \varphi^n$ which is the same as $z \mapsto \varphi^n(z)$. It is well known that composition with a Möbius map is a completely isometric automorphism of the disc algebra.

Suppose that there exists $\tilde{\alpha} : \mathbb{Z} \rightarrow \text{Aut}(\mathcal{C})$ such that $\tilde{\alpha}_n(i(f)) = i(\alpha_n(f)), \forall f \in A(\mathbb{D})$. Calculating

$$\begin{aligned} \varphi(z) &= \frac{z - \frac{1}{2}}{1 - \frac{z}{2}} = \left(z - \frac{1}{2}\right) \left(1 + \frac{z}{2} + \frac{z^2}{4} + \dots\right) \\ &= -\frac{1}{2} + \frac{3}{4}z + \frac{3}{8}z^2 + \dots \end{aligned}$$

we get that

$$\iota(\varphi(z)) = \varphi(z) \oplus \begin{bmatrix} -\frac{1}{2} & 0 \\ \frac{3}{4} & -\frac{1}{2} \end{bmatrix}.$$

Hence,

$$\begin{aligned}
\tilde{\alpha}_1(0 \oplus \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}) &= \tilde{\alpha}_1(\iota(z) - \iota(z^2)\iota(z)^*) \\
&= \iota(\alpha_1(z)) - \iota(\alpha_1(z^2))\iota(\alpha(z))^* \\
&= \iota(\varphi(z)) - \iota(\varphi(z))^2\iota(\varphi(z))^* \\
&= \varphi(z) \oplus \begin{bmatrix} -\frac{1}{2} & 0 \\ \frac{3}{4} & -\frac{1}{2} \end{bmatrix} - \left(\varphi(z)^2 \oplus \begin{bmatrix} \frac{1}{4} & 0 \\ -\frac{3}{4} & \frac{1}{4} \end{bmatrix} \right) \left(\overline{\varphi(z)} \oplus \begin{bmatrix} -\frac{1}{2} & \frac{3}{4} \\ 0 & -\frac{1}{2} \end{bmatrix} \right) \\
&= 0 \oplus \begin{bmatrix} -\frac{3}{8} & -\frac{3}{16} \\ \frac{3}{8} & \frac{3}{16} \end{bmatrix}.
\end{aligned}$$

But then

$$\left\| 0 \oplus \begin{bmatrix} -\frac{3}{8} & -\frac{3}{16} \\ \frac{3}{8} & \frac{3}{16} \end{bmatrix} \right\| = \sqrt{2} \left\| \begin{bmatrix} \frac{3}{8} \\ \frac{3}{16} \end{bmatrix} \right\| = \frac{3\sqrt{10}}{16} < 1 = \left\| 0 \oplus \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \right\|,$$

a contradiction as $*$ -automorphisms are isometric. Therefore, no such $\tilde{\alpha}$ exists and (\mathcal{C}, ι) is a non α -admissible C^* -cover of $(A(\mathbb{D}), \alpha, \mathbb{Z})$. \blacksquare

If we do have an α -admissible cover then we can abuse the notation and call the group representation α again because of the next result.

Lemma 2.2. *Let $(\mathcal{A}, \mathcal{G}, \alpha)$ be an operator algebra dynamical system. If (\mathcal{C}, ι) is an α -admissible C^* -cover then there is a unique group representation of \mathcal{G} on \mathcal{C} acting by $*$ -automorphisms extending α .*

Proof. Let β_1 and β_2 be two such extensions. Then $\beta_1 \circ \beta_2^{-1} = \text{id}$ on \mathcal{A} and we just need to prove this is the identity map. To this end assume that $\beta : \mathcal{G} \rightarrow \text{Aut}(\mathcal{C})$ extends the identity map on \mathcal{A} . That is, $\beta(g)|_{\mathcal{A}} = \text{id}_{\mathcal{A}}$, for all $g \in \mathcal{G}$.

By the universal property of $C_{\max}^*(\mathcal{A})$ there is a unique surjective $*$ -homomorphism $\varphi : C_{\max}^*(\mathcal{A}) \rightarrow \mathcal{C}$ such that $\varphi(a) = \iota(a)$, for all $a \in \mathcal{A}$. Notice that $\beta \circ \varphi(a) = \beta \circ \iota(a) = \iota(a)$, for all $a \in \mathcal{A}$. Thus, $\beta \circ \varphi = \varphi$ by uniqueness which implies that $\beta = \text{id}$ on \mathcal{C} . \blacksquare

Now we turn to crossed products of non-selfadjoint operator algebras.

Definition 2.3 ([20]). Let $(\mathcal{A}, \mathcal{G}, \alpha)$ be an operator algebra dynamical system and let (\mathcal{C}, ι) be an α -admissible C^* -cover. The *relative reduced and full crossed products* are denoted by $\mathcal{A} \rtimes_{(\mathcal{C}, \iota), \alpha}^r \mathcal{G}$ and $\mathcal{A} \rtimes_{(\mathcal{C}, \iota), \alpha} \mathcal{G}$ and are defined to be the closure of $C_c(\mathcal{G}, \mathcal{A})$ in $\mathcal{C} \rtimes_{\alpha}^r \mathcal{G}$ and $\mathcal{C} \rtimes_{\alpha} \mathcal{G}$, respectively.

All relative reduced crossed products are in fact completely isometrically isomorphic [20, Theorem 3.12] and so we define the *reduced crossed product*, denoted $\mathcal{A} \rtimes_{\alpha}^r \mathcal{G}$, to be this unique object. Lastly, we define the *full crossed product* to be

$$\mathcal{A} \rtimes_{\alpha} \mathcal{G} := \mathcal{A} \rtimes_{C_{\max}^*(\mathcal{A}), \alpha} \mathcal{G}.$$

In fact, this is the universal algebra for all covariant representations of $(\mathcal{A}, \mathcal{G}, \alpha)$ [20, Proposition 3.7]. Finally, it should be noted that, as in the selfadjoint case, if \mathcal{G} is amenable then the full and reduced crossed products coincide [20, Theorem 3.14].

Now we are able to state and prove the main theorem of this section.

Theorem 2.4. *Let $(\mathcal{A}, \mathcal{G}, \alpha)$ be an operator algebra dynamical system. Then for every α -admissible C^* -cover (\mathcal{C}, ι) there are surjective completely contractive homomorphisms*

$$\mathcal{A} \rtimes_{\alpha} \mathcal{G} \xrightarrow{q_{\max}} \mathcal{A} \rtimes_{(\mathcal{C}, \iota), \alpha} \mathcal{G} \xrightarrow{q_{\min}} \mathcal{A} \rtimes_{C_{\text{env}}^*(\mathcal{A}), \alpha}$$

such that they are just the identity on $C_c(\mathcal{G}, \mathcal{A})$.

Proof. Label the unique extensions of α to the C^* -covers $C_{\text{env}}^*(\mathcal{A})$, (\mathcal{C}, ι) and $C_{\text{max}}^*(\mathcal{A})$ respectively α_{env} , $\alpha_{\mathcal{C}}$ and α_{max} . By the universal properties there exists surjective $*$ -homomorphisms

$$\varphi_{\text{env}} : \mathcal{C} \rightarrow C_{\text{env}}^*(\mathcal{A}) \quad \text{and} \quad \varphi_{\text{max}} : C_{\text{max}}^*(\mathcal{A}) \rightarrow \mathcal{C}$$

such that $\varphi_{\text{env}}(\iota(a)) = a$ and $\varphi_{\text{max}}(a) = \iota(a)$, for all $a \in \mathcal{A}$.

By uniqueness of the quotient maps and of the extensions we have the following commutative diagram:

$$\begin{array}{ccccc} C_{\text{max}}^*(\mathcal{A}) & \xrightarrow{\varphi_{\text{max}}} & \mathcal{C} & \xrightarrow{\varphi_{\text{env}}} & C_{\text{env}}^*(\mathcal{A}) \\ \alpha_{\text{max}} \downarrow & & \alpha_{\mathcal{C}} \downarrow & & \alpha_{\text{env}} \downarrow \\ C_{\text{max}}^*(\mathcal{A}) & \xrightarrow{\varphi_{\text{max}}} & \mathcal{C} & \xrightarrow{\varphi_{\text{env}}} & C_{\text{env}}^*(\mathcal{A}) \end{array}$$

Thus, $\ker \varphi_{\text{max}}$ is an α_{max} -invariant ideal and $\ker \varphi_{\text{env}}$ is an $\alpha_{\mathcal{C}}$ -invariant ideal. By [32, Proposition 3.19], full C^* -crossed products preserve exact sequences by α -invariant ideals. Hence, we have the following surjective $*$ -homomorphisms

$$C_{\text{max}}^*(\mathcal{A}) \rtimes_{\alpha_{\text{max}}} \mathcal{G} \xrightarrow{\varphi_{\text{max}} \rtimes \text{id}} \mathcal{C} \rtimes_{\alpha_{\mathcal{C}}} \mathcal{G} \xrightarrow{\varphi_{\text{env}} \rtimes \text{id}} C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha_{\text{env}}} \mathcal{G}.$$

So

$$q_{\max} = \varphi_{\text{max}} \rtimes \text{id}|_{\mathcal{A} \rtimes_{\alpha} \mathcal{G}} \quad \text{and} \quad q_{\min} = \varphi_{\text{env}} \rtimes \text{id}|_{\mathcal{A} \rtimes_{(\mathcal{C}, \iota), \alpha} \mathcal{G}}$$

are completely contractive homomorphisms which amount to the identity on $C_c(\mathcal{G}, \mathcal{A})$. ■

The benefit of this theorem, as will be used later, is that one needs only to show that the map $q_{\min} \circ q_{\max}$ is a completely isometric isomorphism to establish that all relative crossed products are the same.

3. HYPERRIGIDITY AND THE HAO-NG ISOMORPHISM

A not necessarily unital operator algebra \mathcal{A} is said to be *hyperrigid* if given any (non-degenerate) $*$ -homomorphism

$$\tau: C_{\text{env}}^*(\mathcal{A}) \longrightarrow B(\mathcal{H})$$

then τ is the only completely positive, completely contractive extension of the restricted map $\tau|_{\mathcal{A}}$. By adding an injective direct summand if necessary, it is easy to see that in order to verify hyperrigidity, one needs to consider only injective $*$ -representations τ but this need not concern us here. The term hyperrigid was coined by Arveson in [2] but the concept had been floating around in various forms before this, e.g. [9].

Our definition is slightly weaker than that of Duncan's [9, Section 4] as Duncan requests that τ be the only completely contractive extension of the restricted map, i.e., no requirement of positivity in the non-unital case. In any case [9, Proposition 4] shows that the graph algebra of any row-finite graph is hyperrigid. Actually we are about to provide a much stronger result but first we need to remind the reader the definition and some of the basic notation regarding C^* -correspondences.

A C^* -correspondence $(X, \mathcal{C}, \varphi_X)$ (or just (X, \mathcal{C})) consists of a C^* -algebra \mathcal{C} , a Hilbert \mathcal{C} -module $(X, \langle \cdot, \cdot \rangle)$ and a (non-degenerate) $*$ -homomorphism $\varphi_X: \mathcal{C} \rightarrow \mathcal{L}(X)$ into the C^* -algebra of adjointable operators on X .

A representation (ρ, t) of a C^* -correspondence into $B(\mathcal{H})$, is a pair consisting of a non-degenerate $*$ -homomorphism $\rho: \mathcal{C} \rightarrow B(\mathcal{H})$ and a linear map $t: X \rightarrow B(\mathcal{H})$, such that

$$\rho(c)t(x) = t(\varphi_X(c)(x)),$$

for all $c \in \mathcal{C}$ and $x \in X$. It is called completely contractive when t is a completely contractive map from the norm induced by the inner product to the operator norm on $B(\mathcal{H})$. Moreover, it is called an isometric (Toeplitz) representation when

$$t(x)^*t(x') = \rho(\langle x, x' \rangle),$$

for all $c \in \mathcal{C}$ and $x, x' \in X$. Note that an isometric representation is necessarily completely contractive but not the other way around.

By the relations above, the C^* -algebra generated by an isometric representation (ρ, t) equals the closed linear span of

$$t(x_1) \cdots t(x_n)t(y_1)^* \cdots t(y_m)^*, \quad x_i, y_j \in X.$$

For any isometric representation (ρ, t) there exists a $*$ -homomorphism $\psi_t : \mathcal{K}(X) \rightarrow B$, such that $\psi_t(\theta_{x,y}) = t(x)t(y)^*$, where $\mathcal{K}(X)$ is the subalgebra of $\mathcal{L}(X)$ of so-called compact operators generated by $\theta_{x,y}(z) = x\langle y, z \rangle$.

There exists a universal Toeplitz representation, denoted as (ρ_∞, t_∞) , so that any other representation of (X, \mathcal{C}) is equivalent to a direct sum of sub-representations of (ρ_∞, t_∞) . The Cuntz-Pimsner-Toeplitz C^* -algebra \mathcal{T}_X is defined as the C^* -algebra generated by the image of (ρ_∞, t_∞) .

The *tensor algebra* \mathcal{T}_X^+ of a C^* -correspondence [26] (X, \mathcal{C}) is the norm-closed subalgebra of \mathcal{T}_X generated by all elements of the form $\rho_\infty(c), t_\infty(x)$, $c \in \mathcal{C}$, $x \in X$.

Consider the ideal

$$\mathcal{J}_X \equiv \ker \varphi_X^\perp \cap \varphi_X^{-1}(\mathcal{K}(X)).$$

(which we will call Katsura's ideal.) An isometric representation (ρ, t) of $(X, \mathcal{C}, \varphi_X)$ is said to be *covariant (Cuntz-Pimsner)* if and only if $\psi_t(\varphi_X(c)) = \rho(c)$, for all $c \in \mathcal{J}_X$. The universal C^* -algebra for all isometric covariant representations of (X, \mathcal{C}) is the Cuntz-Pimsner algebra \mathcal{O}_X . The algebra \mathcal{O}_X contains (a faithful copy of) \mathcal{C} and (a unitarily equivalent) copy of X .

The first author and Kribs [19, Lemma 3.5] have shown that the non-selfadjoint algebra of \mathcal{O}_X generated by these copies of \mathcal{C} and X is completely isometrically isomorphic to \mathcal{T}_X^+ . Furthermore, $C_{\text{env}}^*(\mathcal{T}_X^+) \simeq \mathcal{O}_X$. See [19, 26] for more details.

Now to the hyperrigidity of tensor algebras.

Theorem 3.1. *Let (X, \mathcal{C}) be a C^* -correspondence with X countably generated as a right Hilbert \mathcal{C} -module. If $\varphi_X(\mathcal{J}_X)$ acts non-degenerately on X , then (X, \mathcal{C}) is a hyperrigid C^* -correspondence, i.e., \mathcal{T}_X^+ is a hyperrigid operator algebra.*

Proof. Let $\tau : \mathcal{O}_X \rightarrow B(\mathcal{H})$ be a $*$ -homomorphism and let $\tau' : \mathcal{O}_X \rightarrow B(\mathcal{H})$ be a completely contractive and completely positive map that agrees with τ on \mathcal{T}_X^+ . We are to prove that τ' is multiplicative and so it agrees with τ . Since τ' is a completely contractive and completely positive map, we can use multiplicative domain arguments [6, Proposition 1.5.7].

Let (ρ, t) be the universal Cuntz-Pimsner representation of (X, \mathcal{C}) . Since $\rho(\mathcal{C}) \subseteq \mathcal{T}_X^+$ is a C^* -algebra, the multiplicative domain of τ' contains $\rho(\mathcal{C})$. We claim that it also contains $t(X)$.

Indeed, for any $x \in X$ we have

$$\begin{aligned}
 \tau'(t(x))^* \tau'(t(x)) &= \tau(t(x))^* \tau(t(x)) = \tau(t(x)^* t(x)) \\
 (3) \qquad \qquad \qquad &= \tau(\rho(\langle x, x \rangle)) = \tau'(\rho(\langle x, x \rangle)) \\
 &= \tau'(t(x)^* t(x)),
 \end{aligned}$$

where the equation on the second line holds because $\rho(\mathcal{C}) \subseteq \mathcal{T}_X^+$ and the two maps agree there.

Since X is countably generated, Kasparov's Stabilization Theorem implies the existence of a sequence $\{x_n\}_{n=1}^\infty$ in X so that $\sum_{n=1}^k \theta_{x_n, x_n}$, $k = 1, 2, \dots$, is an approximate unit for $\mathcal{K}(X)$. Let $a \in \mathcal{J}_X$. Then

$$\begin{aligned}
 \varphi_X(aa^*) &= \lim_k \varphi_X(a) \left(\sum_{n=1}^k \theta_{x_n, x_n} \right) \varphi_X(a)^* \\
 &= \sum_{n=1}^\infty \theta_{\varphi_X(a)x_n, \varphi_X(a)x_n},
 \end{aligned}$$

with the convergence in the norm topology. By the Schwarz inequality

$$(4) \quad \tau'(t(\varphi_X(a)x_n)t(\varphi_X(a)x_n)^*) \geq \tau'(t(\varphi_X(a)x_n))\tau'(t(\varphi_X(a)x_n))^*,$$

for all $n \in \mathbb{N}$, and so

$$\begin{aligned}
 \tau'(\rho(aa^*)) &= \tau'(\psi_t(\varphi_X(aa^*))) \\
 &= \sum_{n=1}^\infty \tau'(t(\varphi_X(a)x_n)t(\varphi_X(a)x_n)^*) \\
 &\geq \sum_{n=1}^\infty \tau'(t(\varphi_X(a)x_n))\tau'(t(\varphi_X(a)x_n))^* \\
 &= \sum_{n=1}^\infty \tau(t(\varphi_X(a)x_n))\tau(t(\varphi_X(a)x_n))^* \\
 &= \tau(\psi_t(\varphi_X(aa^*))) = \tau(\rho(aa^*)) = \tau'(\rho(aa^*)).
 \end{aligned}$$

Hence (4) is actually an equality. Combining this with (3), we conclude that $t(\varphi_X(a)x_n)$ belongs to the multiplicative domain of τ' , for all $a \in \mathcal{J}_X$ and $n \in \mathbb{N}$. Since $\rho(\mathcal{C})$ is also contained in the multiplicative domain of τ' , we have that

$$t(\varphi_X(a)x) = \sum_{n=1}^\infty t(\varphi_X(a)x_n)\rho(\langle x_n, x \rangle)$$

belongs to the multiplicative domain of τ' , for all $a \in \mathcal{J}_X$ and $x \in X$. Since $\varphi_X(\mathcal{J}_X)$ acts non-degenerately on X , the multiplicative domain of τ' contains $t(X)$, as desired. This completes the proof. \blacksquare

Recall that a C^* -correspondence (X, \mathcal{C}) is said to be *regular* iff \mathcal{C} acts faithfully on X by compact operators, i.e., $\mathcal{J}_X = \mathcal{C}$. The following is immediate.

Corollary 3.2. *A regular, countably generated C^* -correspondence is necessarily hyperrigid.*

We are about to see that the assumption of injectivity cannot be removed from the Corollary above. But first we need criterion for the failure of hyperrigidity.

Proposition 3.3. *Let (X, \mathcal{C}) be a C^* -correspondence with $\mathcal{J}_X = \{0\}$. Then (X, \mathcal{C}) fails to be hyperrigid.*

Proof. Let (π, t) be any isometric representation of (X, \mathcal{C}) on a Hilbert space \mathcal{H} . If V_1, V_2 are the unilateral and bilateral (forward) shift respectively, then the associations

$$(5) \quad \begin{aligned} \mathcal{C} \ni a &\longrightarrow a \otimes I, \\ X \ni x &\longrightarrow x \otimes V_i, \quad i = 1, 2, \end{aligned}$$

determine isometric representations of X , which are necessarily Cuntz-Pimsner covariant, since $\mathcal{J}_X = \{0\}$. Therefore they promote to representations φ_1 and φ_2 of $\mathcal{O}_X \simeq C_{\text{env}}^*(\mathcal{T}_X^+)$ on $\mathcal{H} \otimes \ell^2(\mathbb{Z}^+)$ and $\mathcal{H} \otimes \ell^2(\mathbb{Z})$ respectively. Now notice that when φ_2 is being compressed on $\mathcal{H} \otimes \ell^2(\mathbb{Z}^+)$, it produces a completely positive contractive map $\tilde{\varphi}_2 \neq \varphi_1$, which however agrees with φ_1 on \mathcal{T}_X^+ . Hence (X, \mathcal{C}) is not hyperrigid. ■

Recall that if (\mathcal{A}, α) is a (non-degenerate) C^* -dynamical system, then the semicrossed product $\mathcal{A} \rtimes_{\alpha} \mathbb{Z}^+$ is simply the tensor algebra of the C^* -correspondence \mathcal{A}_{α} , where the left action on \mathcal{A} is coming from α . In the case where both \mathcal{A} and α are unital and α is injective, such algebras are always hyperrigid. This has already been noted in the literature, eg. [15], but it is also an immediate consequence of Corollary 3.2. It is worth noting that the requirement of α being injective cannot be dropped from neither Corollary 3.2 nor the discussion above.

Example 3.4. Let \mathcal{X} be a connected, compact Hausdorff space and $x \in \mathcal{X}$. Let $\varphi : \mathcal{X} \rightarrow \mathcal{X}$ with $\varphi(y) = x$, for all $y \in \mathcal{X}$. Then the semicrossed product $C(\mathcal{X}) \rtimes_{\varphi} \mathbb{Z}^+$ is not hyperrigid.

Indeed, in that case, the kernel of the right action equals $C_0(\mathcal{X} \setminus \{x\})$. Hence Katsura's ideal is trivial and Proposition 3.3 applies.

Finally recall that the C^* -envelope of a non-unital operator algebra can be computed from the C^* -envelope of its unitization. More precisely, as the pair $(C_{\text{env}}^*(\mathcal{A}), \iota)$ where $C_{\text{env}}^*(\mathcal{A})$ is the C^* -subalgebra generated by $\iota(\mathcal{A})$ inside the C^* -envelope $(C_{\text{env}}^*(\mathcal{A}^1), \iota)$ of the (unique)

unitization \mathcal{A}^1 of \mathcal{A} . By the proof of [5, Proposition 4.3.5] this C^* -envelope of an operator algebra \mathcal{A} has the desired universal property, that for any C^* -cover (ι', \mathcal{B}') of \mathcal{A} , there exists a (necessarily unique and surjective) $*$ -homomorphism $\pi : \mathcal{B}' \rightarrow C_{\text{env}}^*(\mathcal{A})$, such that $\pi \circ \iota' = \iota$.

We start with an elementary result regarding crossed products.

Lemma 3.5. *Let $(\mathcal{C}, \mathcal{G}, \alpha)$ be a C^* -dynamical system and let $\mathcal{D} \subseteq \mathcal{C}$ be the C^* -subalgebra of \mathcal{C} generated by some selfadjoint approximate unit for \mathcal{C} . Then*

- (i) $\mathcal{C}C_c(\mathcal{G}, \mathcal{D})$ is dense in $\mathcal{C} \rtimes_{\alpha} \mathcal{G}$.
- (ii) If $\pi : \mathcal{C} \rtimes_{\alpha} \mathcal{G} \rightarrow B(\mathcal{H})$ is a non-degenerate representation, then its restriction on $C_c(\mathcal{G}, \mathcal{D})$ is also non-degenerate.

Proof. Let $\{e_i\}_{i \in \mathbb{I}}$ be the selfadjoint approximate unit generating \mathcal{D} . Then any elementary tensor $h \otimes c \in C_c(\mathcal{G}, \mathcal{D})$, where $(h \otimes c)(s) = h(s)c$, $h \in C_c(\mathcal{G})$, $c \in \mathcal{C}$, can be written as

$$h \otimes c = \lim_{i \in \mathbb{I}} c(h \otimes e_i) \in \overline{\mathcal{C}C_c(\mathcal{G}, \mathcal{D})}.$$

This implies (i). For (ii) notice that by taking adjoints in (i), $C_c(\mathcal{G}, \mathcal{D})\mathcal{C}$ is also dense in $\mathcal{C} \rtimes_{\alpha} \mathcal{G}$. Hence

$$\pi(C_c(\mathcal{G}, \mathcal{D}))\mathcal{H} = \pi(C_c(\mathcal{G}, \mathcal{D}))\pi(\mathcal{C})\mathcal{H} = \pi(C_c(\mathcal{G}, \mathcal{D})\mathcal{C})\mathcal{H}$$

which is dense in $\pi(\mathcal{C})\mathcal{H} = \mathcal{H}$ and the conclusion follows. \blacksquare

The following result has been established for all discrete groups in [18]. Here we extend it to arbitrary locally compact groups provided that the pertinent algebras are hyperrigid.

Theorem 3.6. *Let \mathcal{A} be a hyperrigid operator algebra which possesses a contractive approximate unit $\{e_i\}_{i \in \mathbb{I}}$ consisting of selfadjoint operators. Let $\alpha : \mathcal{G} \rightarrow \text{Aut } \mathcal{A}$ be a continuous action of a locally compact group. Then*

$$(6) \quad C_{\text{env}}^*(\mathcal{A} \rtimes_{\alpha}^r \mathcal{G}) \simeq C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G}$$

and

$$(7) \quad C_{\text{env}}^*(\mathcal{A} \rtimes_{C_{\text{env}}^*(\mathcal{A}), \alpha} \mathcal{G}) \simeq C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha} \mathcal{G}$$

via canonical embeddings.

Proof. Let $\rho : C_{\text{env}}^*(\mathcal{A}) \rightarrow B(\mathcal{H})$ be a faithful (non-degenerate) representation and let

$$\begin{aligned} \tilde{\rho} : C_{\text{env}}^*(\mathcal{A}) &\longrightarrow B(\mathcal{H} \otimes L^2(\mathcal{G})) \\ u : \mathcal{G} &\longrightarrow B(\mathcal{H} \otimes L^2(\mathcal{G})) \end{aligned}$$

so that $\tilde{\rho} \rtimes u$ (which we will denote as π) is the regular representation induced by ρ . (See [32, Section 2.2] for notation and additional information.) Since ρ is non-degenerate, [32, Lemma 2.17] implies that the induced representation $\pi = \tilde{\rho} \otimes u$ is also non-degenerate.

Let

$$\varphi: B(\mathcal{H} \otimes L^2(\mathcal{G})) \longrightarrow B(\mathcal{H} \otimes L^2(\mathcal{G}))$$

be a contractive idempotent map whose range is the injective envelope of $\pi(\mathcal{A} \rtimes_{\alpha}^r \mathcal{G})^1$. Let \mathcal{D} be the closed (selfadjoint) subalgebra of \mathcal{A} generated by $\{e_i\}_{i \in \mathbb{I}}$. Then $C_c(\mathcal{G}, \mathcal{D})$ is a selfadjoint subalgebra of $\mathcal{A} \rtimes_{\alpha}^r \mathcal{G}$ and so [5, 1.3.12] implies that

$$(8) \quad \varphi(S\pi(f)) = \varphi(S)\pi(f),$$

for any $S \in B(\mathcal{H} \otimes L^2(\mathcal{G}))$ and $f \in C_c(\mathcal{G}, \mathcal{D})$. In particular

$$\varphi(\tilde{\rho}(a)\pi(f)) = \varphi(\tilde{\rho}(a))\pi(f),$$

for all $a \in \mathcal{A}$, $f \in C_c(\mathcal{G}, \mathcal{D})$. On the other hand,

$$\tilde{\rho}(a)\pi(f) = \pi(af) \in \pi(\mathcal{A} \rtimes_{\alpha}^r \mathcal{G})$$

and so $\varphi(\tilde{\rho}(a)\pi(f)) = \tilde{\rho}(a)\pi(f)$, $a \in \mathcal{A}$, $f \in C_c(\mathcal{G}, \mathcal{D})$. Hence

$$(\varphi(\tilde{\rho}(a)) - \tilde{\rho}(a))\pi(f) = 0, \text{ for all } f \in C_c(\mathcal{G}, \mathcal{D}).$$

By Lemma 3.5(ii), $\pi(C_c(\mathcal{G}, \mathcal{D}))$ acts non-degenerately on $\mathcal{H} \otimes L^2(\mathcal{G})$ and so

$$(9) \quad \varphi(\tilde{\rho}(a)) = \tilde{\rho}(a), \text{ for all } a \in \mathcal{A}.$$

Hence the mapping φ is a completely positive and completely contractive extension of the identity map on $\tilde{\rho}(\mathcal{A})$. However $\tilde{\rho}(\mathcal{A})$ is hyper-rigid and according to the discussion in the beginning of the section, the identity map on $\tilde{\rho}(\mathcal{A})$ is the only such completely contractive and completely positive extension to $C_{\text{env}}^*(\tilde{\rho}(\mathcal{A})) = \tilde{\rho}(C_{\text{env}}^*(\mathcal{A}))$. Therefore

$$\varphi(\tilde{\rho}(c)) = \tilde{\rho}(c), \text{ for all } c \in C_{\text{env}}^*(\mathcal{A}).$$

Appealing again to (8), with $S = \tilde{\rho}(c)$, we obtain

$$\varphi(\pi(cf)) = \varphi(\tilde{\rho}(c)\pi(f)) = \varphi(\tilde{\rho}(c))\pi(f) = \tilde{\rho}(c)\pi(f) = \pi(cf),$$

for all $c \in C_{\text{env}}^*(\mathcal{A})$ and $f \in C_c(\mathcal{G}, \mathcal{D})$. By Lemma 3.5(i) we have

$$\varphi(S) = S, \text{ for all } S \in \pi(C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G}).$$

But this implies that the Choi-Effros product on

$$\pi(C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G})^1 \subseteq \varphi(B(\mathcal{H} \otimes L^2(\mathcal{G})))$$

is actually the one coming from $B(\mathcal{H} \otimes L^2(\mathcal{G}))$ and the C^* -algebra generated by

$$\pi(\mathcal{A} \rtimes_{\alpha}^r \mathcal{G})^1 \subseteq \varphi(B(\mathcal{H} \otimes L^2(\mathcal{G})))$$

equals $\pi(C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G})^1$. Hence

$$C_{\text{env}}^*(\pi(\mathcal{A} \rtimes_{\alpha}^r \mathcal{G})^1) = \pi(C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G})^1.$$

Furthermore, the C^* -algebra generated by $\pi(\mathcal{A} \rtimes_{\alpha}^r \mathcal{G}) \subseteq \pi(C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G})^1$ equals $\pi(C_{\text{env}}^*(\mathcal{A}) \rtimes_{\alpha}^r \mathcal{G})$. This establishes (6).

In order to prove (7), let this time $\pi := \tilde{\rho} \rtimes u$, where $(\tilde{\rho}, u)$ is the universal covariant representation of $(C_{\text{env}}^*(\mathcal{A}), \mathcal{G}, \alpha)$. With this π , a verbatim repetition of the proof of (6) establishes (7). \blacksquare

Now we turn to crossed product correspondences. Let \mathcal{G} be a locally compact group acting on a non-degenerate C^* -correspondence (X, \mathcal{C}) by a generalized gauge action $\alpha : \mathcal{G} \rightarrow \text{Aut}(\mathcal{T}_X)$. The reduced crossed product correspondence $(X \rtimes_{\alpha}^r \mathcal{G}, \mathcal{C} \rtimes_{\alpha}^r \mathcal{G})$ is the completion of $C_c(\mathcal{G}, X)$ and $C_c(\mathcal{G}, \mathcal{C})$ in $\mathcal{T}_X^+ \rtimes_{\alpha}^r \mathcal{G}$, which is usually thought of as living in $\mathcal{O}_X \rtimes_{\alpha}^r \mathcal{G}$ but equivalently can be considered as living in $\mathcal{T}_X \rtimes_{\alpha}^r \mathcal{G}$. The left and right module actions are given by multiplication and $\langle S, T \rangle = S^*T$, for $S, T \in C_c(\mathcal{G}, X)$.

In a similar manner, one defines the full crossed product correspondence $(X \rtimes_{\alpha} \mathcal{G}, \mathcal{C} \rtimes_{\alpha} \mathcal{G})$ by completing the spaces in $\mathcal{T}_X \rtimes_{\alpha} \mathcal{G}$. This was shown to be unitarily equivalent to the abstract characterization of the full crossed product correspondence in [20, Remark 7.8]. Lastly, we recall the definition of the crossed product correspondence $(X \hat{\rtimes}_{\alpha} \mathcal{G}, \mathcal{C} \hat{\rtimes}_{\alpha} \mathcal{G})$ which is the completion of the spaces in $\mathcal{O}_X \rtimes_{\alpha} \mathcal{G}$. In general, it is unknown whether these two correspondences are unitarily equivalent or not.

The following has been established in [20] in the case where \mathcal{G} is discrete. In [20] it was also noted that the proof carries over to the general locally compact case. We now explain how this is done.

Theorem 3.7. *Let \mathcal{G} be a locally compact group acting on a non-degenerate C^* -correspondence (X, \mathcal{C}) . Then*

$$\mathcal{T}_X^+ \rtimes_{\alpha}^r \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha}^r \mathcal{G}}^+.$$

Therefore,

$$C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\alpha}^r \mathcal{G}) \simeq \mathcal{O}_{X \rtimes_{\alpha}^r \mathcal{G}}.$$

Proof. Let $\rho : \mathcal{T}_X \rightarrow B(\mathcal{H})$ be some faithful $*$ -representation. Let \mathbb{Z}^+ denote the non-negative integers and let $V \in B(\ell^2(\mathbb{Z}^+))$ be the forward shift. Set $X_n := X^{\otimes n}$ for $n \geq 1$, $X_0 := \mathcal{C}$ and for the rest of the proof

let X denote the product system $\{X_n\}_{n=0}^\infty$. (See [11] for the definition and notation on product systems.) Then the map

$$X_n \ni x \longmapsto \rho(x) \otimes V^n \in B(\mathcal{H} \otimes \ell^2(\mathbb{Z}^+)), \quad n \in \mathbb{Z}^+,$$

is a Toeplitz representation i.e., Nica-covariant representation, of the product system X that satisfies the requirements of Fowler's Theorem [11, Theorem 7.2]. Therefore it establishes a faithful representation $\pi : \mathcal{T}_X \rightarrow B(\mathcal{H} \otimes \ell^2(\mathbb{Z}^+))$.

Since $X \rtimes_\alpha^r \mathcal{G} \subseteq \mathcal{T}_X \rtimes_\alpha^r \mathcal{G}$, we may consider the regular representation Ind_π , when restricted on $X \rtimes_\alpha^r \mathcal{G}$, as a Toeplitz representation of the product system $X \rtimes_\alpha^r \mathcal{G}$, which we denote as ψ . We furthermore write $\psi_n := \psi|_{X_n \rtimes_\alpha^r \mathcal{G}}$, $n \in \mathbb{Z}^+$.

We claim that ψ satisfies the requirements of Fowler's Theorem [11, Theorem 7.2]. Indeed let $Q_n \in \ell^2(\mathbb{Z}^+)$ be the projection on the one dimensional subspace corresponding to the characteristic function of $n \in \mathbb{Z}^+$ and let $\hat{Q}_n \equiv (I \otimes Q_n) \otimes I \in B(\mathcal{H} \otimes \ell^2(\mathbb{Z}^+) \otimes L^2(\mathcal{G}))$ be the (constant) $B(\mathcal{H} \otimes \ell^2(\mathbb{Z}^+))$ -valued function that assigns the value $I \otimes Q_n$ to any $s \in \mathcal{G}$. Then given any $f \in C_c(\mathcal{G}, X_n)$, $n \geq 1$, we have

$$\begin{aligned} (\hat{Q}_0 \text{Ind}_\pi f)h(t) &= \hat{Q}_0 \int_{\mathcal{G}} \pi(\alpha_t^{-1}(f(s)))h(s^{-1}t)ds \\ &= (I \otimes Q_0) \otimes I \int_{\mathcal{G}} (\rho(\alpha_t^{-1}(f(s))) \otimes V^n)h(s^{-1}t)ds \\ &= \int_{\mathcal{G}} (\rho(\alpha_t^{-1}(f(s))) \otimes Q_0 V^n)h(s^{-1}t)ds = 0 \end{aligned}$$

Therefore if P_n^ψ denotes the range space of $\psi(X_n \rtimes_\alpha^r \mathcal{G})$, $n \in \mathbb{Z}^+$, then the product $\prod_{n \in F \setminus \{0\}} (I - P_n^\psi)$, which in our case collapses to a single factor of the form $I - P_n^\psi$, always dominates \hat{Q}_0 and so

$$(10) \quad \|\psi_0(f) \prod_{n \in F \setminus \{0\}} (I - P_n^\psi)\| \geq \|\psi_0(f)\hat{Q}_0\| = \|\text{Ind}_\pi(f)\hat{Q}_0\|,$$

for any $f \in C_c(\mathcal{G}, \mathcal{C})$. However, each $I \otimes Q_n$ reduces $\pi|_{\mathcal{C}}$ and therefore

$$\pi|_{\mathcal{C}} \simeq \left(\bigoplus_{n \in \mathbb{Z}^+} (I \otimes Q_n)\pi \right)|_{\mathcal{C}} \simeq \left(\bigoplus (I \otimes Q_0)\pi \right)|_{\mathcal{C}},$$

i.e., the restriction of π on \mathcal{C} is unitarily equivalent to a direct sum of countably many copies of $(I \otimes Q_0)\pi$ restricted on \mathcal{C} . From this we obtain,

$$\psi_0 = \text{Ind}_\pi|_{\mathcal{C} \rtimes_\alpha^r \mathcal{G}} \simeq \bigoplus \text{Ind}_{Q_0\pi}|_{\mathcal{C} \rtimes_\alpha^r \mathcal{G}} \simeq \bigoplus \hat{Q}_0 \text{Ind}_\pi|_{\mathcal{C} \rtimes_\alpha^r \mathcal{G}}.$$

Combining the above with (10) we now obtain

$$\|\psi_0(f) \prod_{n \in F \setminus \{0\}} (I - P_n^\psi)\| = \|\text{Ind}_\pi(f)\hat{Q}_0\| = \|\psi_0(f)\|$$

for any $f \in \mathcal{C}_c(\mathcal{G}, \mathcal{C})$, which establishes the claim.

Since the claim is valid, Fowler's Theorem shows now that the induced representation ψ_* is a faithful representation of $\mathcal{T}_{X \rtimes_\alpha^r \mathcal{G}}$. Note now that $\psi_*(\mathcal{T}_{X \rtimes_\alpha^r \mathcal{G}}^+) \simeq \mathcal{T}_{X \rtimes_\alpha^r \mathcal{G}}^+$ is equal to the closed linear span of

$$\psi_*(X \rtimes_\alpha^r \mathcal{G}) = \bigcup_{n \in \mathbb{Z}^+} \psi_n(X_n \rtimes_\alpha^r \mathcal{G}) = \bigcup_{n \in \mathbb{Z}^+} \overline{\text{Ind}_\pi C_c(\mathcal{G}, X_n)}.$$

However, $\mathcal{T}_X^+ \rtimes_\alpha^r \mathcal{G}$ is also isomorphic to the closed linear span of

$$\bigcup_{n \in \mathbb{Z}^+} \overline{\text{Ind}_\pi C_c(\mathcal{G}, X_n)}.$$

Hence, $\mathcal{T}_X^+ \rtimes_\alpha^r \mathcal{G} \simeq \mathcal{T}_{X \rtimes_\alpha^r \mathcal{G}}^+$.

Finally

$$\text{C}_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_\alpha^r \mathcal{G}) \simeq \text{C}_{\text{env}}^*(\mathcal{T}_{X \rtimes_\alpha^r \mathcal{G}}^+) \simeq \mathcal{O}_{X \rtimes_\alpha^r \mathcal{G}}.$$

with the last identification following from [19, Theorem 3.7]. \blacksquare

Remark 3.8. As the reader may have noticed, the proof of the previous result was given in the language of product systems. This was done intentionally because by switching from \mathbb{N} to an arbitrary quasi-lattice ordered group (G, P) , the same exact proof as above establishes the more general result $\mathcal{N}\mathcal{T}_X^+ \rtimes_\alpha^r \mathcal{G} \simeq \mathcal{N}\mathcal{T}_{X \rtimes_\alpha^r \mathcal{G}}^+$, where $\mathcal{N}\mathcal{T}_X^+$ denotes the Nica tensor algebra of X . We are recording this fact here for future reference.

In [18] the Hao-Ng isomorphism problem was resolved for the reduced crossed product and all discrete groups. In the next result we address the case of an arbitrary locally compact group and we resolve the Hao-Ng problem for the reduced crossed product provided that the C^* -correspondence is hyperrigid. Note that our result subsumes an earlier result of Kim [22, Proposition 5.5] which was posted on the arXiv but has not appeared in print.

Theorem 3.9. *Let \mathcal{G} be a locally compact group acting on a non-degenerate hyperrigid C^* -correspondence (X, \mathcal{C}) , e.g. X is countably generated as a right \mathcal{C} -module and $\varphi_X(\mathcal{J}_X)X = X$. Then*

$$\mathcal{O}_X \rtimes_\alpha^r \mathcal{G} \simeq \mathcal{O}_{X \rtimes_\alpha^r \mathcal{G}}.$$

Proof. By Theorem 3.7 we have $C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\alpha}^r \mathcal{G}) \simeq \mathcal{O}_{X \rtimes_{\alpha}^r \mathcal{G}}$. On the other hand, Theorem 3.6 implies that $C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\alpha}^r \mathcal{G}) \simeq C_{\text{env}}^*(\mathcal{T}_X^+) \rtimes_{\alpha}^r \mathcal{G}$. Hence $\mathcal{O}_X \rtimes_{\alpha}^r \mathcal{G} \simeq C_{\text{env}}^*(\mathcal{T}_X^+) \rtimes_{\alpha}^r \mathcal{G} \simeq \mathcal{O}_{X \rtimes_{\alpha}^r \mathcal{G}}$. ■

It is important to us that an analogous results holds for the full crossed product.

Theorem 3.10. *Let \mathcal{G} be a locally compact group acting on a non-degenerate hyperrigid C^* -correspondence (X, \mathcal{C}) . Then*

$$\mathcal{O}_X \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{O}_{X \hat{\rtimes}_{\alpha} \mathcal{G}}.$$

Proof. In [20, Theorem 7.13] we proved that

$$\mathcal{T}_X^+ \rtimes_{\mathcal{O}_X, \alpha} \mathcal{G} \simeq \mathcal{T}_{X \hat{\rtimes}_{\alpha} \mathcal{G}}^+ \quad \text{and} \quad C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\mathcal{O}_X, \alpha} \mathcal{G}) \simeq \mathcal{O}_{X \hat{\rtimes}_{\alpha} \mathcal{G}}.$$

Now (7) in Theorem 3.6 and the above imply that

$$\mathcal{O}_{X \hat{\rtimes}_{\alpha} \mathcal{G}} \simeq C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{C_{\text{env}}^*(\mathcal{T}_X^+), \alpha} \mathcal{G}) \simeq C_{\text{env}}^*(\mathcal{T}_X^+) \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{O}_X \rtimes_{\alpha} \mathcal{G}$$

and the conclusion follows. ■

At this point one might think that the above theorem is the final word regarding the Hao-Ng isomorphism for hyperrigid correspondences. As it turns out, this couldn't be further from the truth. It is indeed the case that we have expressed the crossed product $\mathcal{O}_X \rtimes_{\alpha} \mathcal{G}$ as the Cuntz-Pimsner algebra of a C^* -correspondence, namely $X \hat{\rtimes}_{\alpha} \mathcal{G}$, but this is not the C^* -correspondence that the authors of [3] ask for. Is this a big deal? Most definitely yes, and we devote the next section explaining the reasons why.

4. ISOMETRIC COEXTENSIONS

The goal of this section is to answer Problem 3 in Chapter 8 of [20]: Is $\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G}$ the tensor algebra of some C^* -correspondence? The (affirmative) answer is one of the key ingredients in the proof of Theorem 4.9, one of the central results of the paper.

Definition 4.1. Let (X, \mathcal{C}) be a C^* -correspondence and \mathcal{G} a locally compact group. A *generalized gauge action* $\alpha : \mathcal{G} \rightarrow \text{Aut}((X, \mathcal{C}))$ is a map from \mathcal{G} into the completely isometric module automorphisms. In particular, for each $s \in \mathcal{G}$, α_s is an isometric automorphism of X and a $*$ -automorphism of \mathcal{C} such that

$$\begin{aligned} \alpha_s(\xi c) &= \alpha_s(\xi) \alpha_s(c), \quad \alpha_s(\varphi_X(c) \xi) = \varphi_X(\alpha_s(c)) \alpha_s(\xi) \\ &\text{and } \alpha_s(\langle \xi, \eta \rangle) = \langle \alpha_s(\xi), \alpha_s(\eta) \rangle \end{aligned}$$

for all $\xi, \eta \in X$ and $c \in \mathcal{C}$.

Alternatively, one says that $\alpha : \mathcal{G} \rightarrow \text{Aut } \mathcal{T}_X$ forms a *generalized gauge action of \mathcal{T}_X* if $\alpha_s(X) = X$ and $\alpha_s(\mathcal{C}) = \mathcal{C}$, for all $g \in \mathcal{G}$. The following was observed in [18, pg 5760]. (See also [13, Lemma 2.6] for the analogous result with \mathcal{O}_X).

Proposition 4.2. *Let (X, \mathcal{C}) be a non-degenerate C^* -correspondence and \mathcal{G} a locally compact group. If $\alpha : \mathcal{G} \rightarrow \text{Aut}(\mathcal{T}_X)$ is a generalized gauge action of \mathcal{T}_X then it restricts to a generalized gauge action of (X, \mathcal{C}) . Conversely, a generalized gauge action α of (X, \mathcal{C}) extends uniquely to a generalized gauge action of \mathcal{T}_X .*

The fundamental object of study in this section is the C^* -correspondence dynamical system $((X, \mathcal{C}), \mathcal{G}, \alpha)$ which is given by a non-degenerate C^* -correspondence (X, \mathcal{C}) and a generalized gauge action α of the locally compact group \mathcal{G} acting on (X, \mathcal{C}) .

Definition 4.3. A representation of the C^* -correspondence dynamical system $((X, \mathcal{C}), \mathcal{G}, \alpha)$ is a quadruple $(\rho, t, u, \mathcal{H})$ consisting of a completely contractive representation (ρ, t, \mathcal{H}) of (X, \mathcal{C}) and a strongly continuous unitary representation $u : \mathcal{G} \rightarrow U(\mathcal{H})$ satisfying the covariance relations

$$u(s)t(\xi) = t(\alpha_s(\xi))u(s) \quad \text{and} \quad u(s)\rho(c) = \rho(\alpha_s(c))u(s)$$

for all $s \in \mathcal{G}, \xi \in X$ and $c \in \mathcal{C}$. Moreover, $(\rho, t, u, \mathcal{H})$ is said to be *isometric* if (ρ, t, \mathcal{H}) is an isometric (Toeplitz) representation of (X, \mathcal{C}) .

The following theorem is an extension of [26, Theorem 2.12].

Theorem 4.4. *Let $((X, \mathcal{C}), \mathcal{G}, \alpha)$ be a C^* -correspondence dynamical system. The isometric representations $(\rho, t, u, \mathcal{H})$ of $((X, \mathcal{C}), \mathcal{G}, \alpha)$ are in bijective correspondence with the isometric representations (π, u, \mathcal{H}) of $(\mathcal{T}_X, \mathcal{G}, \alpha)$. Specifically, they are related by $\pi = \rho \rtimes t$.*

Proof. If (π, u, \mathcal{H}) is an isometric representation of $(\mathcal{T}_X, \mathcal{G}, \alpha)$ then [26, Theorem 2.12] proves that there exists an isometric representation (ρ, t, \mathcal{H}) of (X, \mathcal{C}) such that $\rho \rtimes t = \pi$. Proposition 4.2 gives that α is a generalized gauge action of \mathcal{G} on (X, \mathcal{C}) and so the covariance relations between ρ, t and u are automatic. Hence, $(\rho, t, u, \mathcal{H})$ is an isometric representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$.

Conversely, suppose $(\rho, t, u, \mathcal{H})$ is an isometric representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$. Again by [26, Theorem 2.12] this gives that $(\rho \rtimes t, \mathcal{H})$ is an isometric representation of \mathcal{T}_X and by Proposition 4.2 α extends uniquely to a generalized gauge action of \mathcal{G} on \mathcal{T}_X . Because $u(s)(\rho \rtimes t(\cdot))u(s)^*$ and $(\rho \circ \alpha_s) \rtimes (t \circ \alpha_s) = (\rho \rtimes t) \circ \alpha_s$ agree on X and \mathcal{C} then the uniqueness of [26, Theorem 2.12] gives that

$$u(s)\rho \rtimes t(a) = \rho \rtimes t(\alpha_s(a))u(s)$$

for all $s \in \mathcal{G}$ and $a \in \mathcal{T}_X$. Therefore, $(\rho \rtimes t, u, \mathcal{H})$ is an isometric representation of $(\mathcal{T}_X, \mathcal{G}, \alpha)$. \blacksquare

If $(\rho, t, u, \mathcal{H})$ and $(\rho_1, t_1, u_1, \mathcal{H}_1)$ are completely contractive representations of $((X, \mathcal{C}), \mathcal{G}, \alpha)$ we say that the latter is a *dilation* of the former if $\mathcal{H} \subseteq \mathcal{H}_1$ and

- (i) \mathcal{H} reduces π_1 and u_1 with $\pi_1(c)|_{\mathcal{H}} = \pi(c)$, $c \in \mathcal{C}$ and $u_1(g)|_{\mathcal{H}} = u(g)$, $g \in \mathcal{G}$, and
- (ii) \mathcal{H} is a semi-invariant subspace for t_1 and $P_{\mathcal{H}}t_1(\xi)|_{\mathcal{H}} = t(\xi)$, $\xi \in X$.

We call such a dilation an *extension* if \mathcal{H} is an invariant subspace for t_1 and a *coextension* if \mathcal{H} is a coinvariant subspace for t_1 .

The dilation $(\rho_1, t_1, u_1, \mathcal{H}_1)$ is called *minimal* when \mathcal{H}_1 is the smallest reducing subspace for t_1 containing \mathcal{H} .

Now we need a lemma relating to the step-by-step dilation techniques of Muhly and Solel of [26, Section 3]. Their proof is modelled after Popescu's step-by-step dilation technique [28] using the Schaëffer matrix construction [30]. The following is also a slight simplification of the original proof in [26].

To this end suppose $(\rho, t, u, \mathcal{H})$ is a completely contractive representation of the C^* -correspondence dynamical system $((X, \mathcal{C}), \mathcal{G}, \alpha)$. The ultimate goal is to prove that every such representation dilates to an isometric representation.

As in [26], define the Hilbert space $\mathcal{H}^X = \overline{X \otimes_{\rho} \mathcal{H}}^{\langle \cdot, \cdot \rangle}$ where

$$\xi c \otimes h = \xi \otimes \rho(c)h \quad \text{and} \quad \langle \xi \otimes h, \eta \otimes k \rangle = \langle h, \rho(\langle \xi, \eta \rangle)k \rangle$$

for $\xi, \eta \in X$, $c \in \mathcal{C}$ and $h, k \in \mathcal{H}$. As well, define $\sigma^X : X \rightarrow B(\mathcal{H}, \mathcal{H}^X)$ by $\sigma^X(\xi)h = \xi \otimes h$ and $\tilde{t} : \mathcal{H}^X \rightarrow \mathcal{H}$ by $\tilde{t}(\xi \otimes h) = t(\xi)h$. From here one defines the one step dilation to $\mathcal{H}_1 = \mathcal{H} \oplus \mathcal{H}^X$ given by

$$t_1(\xi) = \begin{bmatrix} t(\xi) & 0 \\ (I - \tilde{t}^*\tilde{t})^{1/2}\sigma^X(\xi) & 0 \end{bmatrix}$$

and

$$\rho_1(c) = \begin{bmatrix} \rho(c) & 0 \\ 0 & \tilde{\rho}(c) \end{bmatrix}$$

where $\tilde{\rho} : \mathcal{C} \rightarrow B(\mathcal{H}^X)$ is given by $\tilde{\rho}(c)(\xi \otimes h) = \varphi_X(c)\xi \otimes h$.

Lemma 4.5. *Consider*

$$u_1(s) = \begin{bmatrix} u(s) & 0 \\ 0 & \tilde{u}(s) \end{bmatrix}$$

where $\tilde{u} : G \rightarrow B(\mathcal{H}^X)$ is given by $\tilde{u}(s)(\xi \otimes h) = \alpha_s(\xi) \otimes u(s)h$, which is well-defined. Then $(\rho_1, t_1, u_1, \mathcal{H}_1)$ is a completely contractive representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$ such that

$$t_1(\xi)^* t_1(\eta) = \begin{bmatrix} \rho(\langle \xi, \eta \rangle) & 0 \\ 0 & 0 \end{bmatrix}$$

for all $\xi, \eta \in \mathcal{H}$.

Proof. By [26, Lemma 3.7] (ρ_1, t_1) is a completely contractive representation of (X, \mathcal{C}) on \mathcal{H}_1 which satisfies the last two statements in the lemma.

First note that \tilde{u} is in fact well-defined since it respects the internal \mathcal{C} -modularity of \mathcal{H}^X ,

$$\begin{aligned} \tilde{u}(s)(\xi c \otimes h) &= \alpha_s(\xi c) \otimes u(s)h \\ &= \alpha_s(\xi)\alpha_s(c) \otimes u(s)h \\ &= \alpha_s(\xi) \otimes \rho(\alpha_s(c))u(s)h \\ &= \alpha_s(\xi) \otimes u(s)\rho(c)h \\ &= \tilde{u}(s)(\xi \otimes \rho(c)h), \end{aligned}$$

for all $s \in \mathcal{G}, c \in \mathcal{C}, \xi \in X, h \in \mathcal{H}$.

Additionally, observe that $\tilde{u}(s)$ is unitary since for all $s \in \mathcal{G}, \xi, \eta \in X$ and $h, k \in \mathcal{H}$ we have that

$$\begin{aligned} \langle \tilde{u}(s)(\xi \otimes h), \tilde{u}(s)(\eta \otimes k) \rangle &= \langle \alpha_s(\xi) \otimes u(s)h, \alpha_s(\eta) \otimes u(s)k \rangle \\ &= \langle u(s)h, \rho(\langle \alpha_s(\xi), \alpha_s(\eta) \rangle)u(s)k \rangle \\ &= \langle u(s)h, \rho(\alpha_s(\langle \xi, \eta \rangle))u(s)k \rangle \\ &= \langle u(s)h, u(s)\rho(\langle \xi, \eta \rangle)k \rangle \\ &= \langle h, \rho(\langle \xi, \eta \rangle)k \rangle \\ &= \langle \xi \otimes h, \eta \otimes k \rangle. \end{aligned}$$

Next we need to make the following calculations:

$$\begin{aligned} \sigma^X(\alpha_s(\xi))u(s)h &= \alpha_s(\xi) \otimes u(s)h \\ &= \tilde{u}(s)(\xi \otimes h) \\ &= \tilde{u}(s)\sigma^X(\xi)h \end{aligned}$$

and

$$\begin{aligned}
 \langle \tilde{t}^* \tilde{t} \sigma^X(\alpha_s(\xi)) u(s) h, \eta \otimes k \rangle &= \langle \tilde{t}(\alpha_s(\xi) \otimes u(s) h), \tilde{t}(\eta \otimes k) \rangle \\
 &= \langle t(\alpha_s(\xi)) u(s) h, t(\eta) k \rangle \\
 &= \langle u(s) t(\xi) h, t(\eta) k \rangle \\
 &= \langle t(\xi) h, t(\alpha_{s^{-1}}(\eta)) u(s)^* k \rangle \\
 &= \langle \tilde{t}^* \tilde{t}(\xi \otimes h), \alpha_{s^{-1}}(\eta) \otimes u(s^{-1}) k \rangle \\
 &= \langle \tilde{u}(s) \tilde{t}^* \tilde{t}(\xi \otimes h), \eta \otimes k \rangle \\
 &= \langle \tilde{u}(s) \tilde{t}^* \tilde{t} \sigma^X(\xi) h, \eta \otimes k \rangle.
 \end{aligned}$$

Combining these one gets

$$\begin{aligned}
 \tilde{u}(s)(I - \tilde{t}^* \tilde{t}) \sigma^X(\xi) &= (I - \tilde{t}^* \tilde{t}) \sigma^X(\alpha_s(\xi)) u(s) \\
 &= (I - \tilde{t}^* \tilde{t}) \tilde{u}(s) \sigma^X(\xi).
 \end{aligned}$$

Hence,

$$\tilde{u}(s)(I - \tilde{t}^* \tilde{t}) = (I - \tilde{t}^* \tilde{t}) \tilde{u}(s)$$

and by a standard trick often attributed to Halmos

$$\tilde{u}(s)(I - \tilde{t}^* \tilde{t})^{1/2} = (I - \tilde{t}^* \tilde{t})^{1/2} \tilde{u}(s).$$

Now, we need to establish the covariance relations between (ρ_1, t_1) and u_1 . From the previous paragraph we have that

$$\tilde{u}(s)(I - \tilde{t}^* \tilde{t})^{1/2} \sigma^X(\xi) = (I - \tilde{t}^* \tilde{t})^{1/2} \sigma^X(\alpha_s(\xi)) u(s)$$

and thus $u_1(s) t_1(\xi) = t_1(\alpha_s(\xi)) u_1(s)$.

Second, it is much more straightforward to calculate that

$$\begin{aligned}
 \tilde{u}(s) \tilde{\rho}(c)(\xi \otimes h) &= \tilde{u}(s)(\varphi_X(c) \xi \otimes h) \\
 &= \alpha_s(\varphi_X(c) \xi) \otimes u(s) h \\
 &= \varphi_X(\alpha_s(c)) \alpha_s(\xi) \otimes u(s) h \\
 &= \tilde{\rho}(\alpha_s(c)) \tilde{u}(s)(\xi \otimes h).
 \end{aligned}$$

Therefore, $u_1(s) \rho_1(c) = \rho_1(\alpha_s(c)) u_1(s)$ and the conclusion follows. \blacksquare

Theorem 4.6. *Every completely contractive representation of the non-degenerate C^* -correspondence dynamical system $((X, \mathcal{C}), \mathcal{G}, \alpha)$ has a minimal isometric coextension. Moreover, the minimal isometric coextension is unique up to unitary equivalence.*

Proof. Let $(\rho, t, u, \mathcal{H})$ be a completely contractive representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$. Following the proof of [26, Theorem 3.3] repeatedly use Lemma 4.5 to get a sequence of completely contractive representations

$(\rho_n, t_n, u_n, \mathcal{H}_n)$ of $((X, \mathcal{C}), \mathcal{G}, \alpha)$ in the obvious manner: use $(\rho, t, u, \mathcal{H})$ to produce $(\rho_1, t_1, u_1, \mathcal{H}_1)$ and then recursively

$$(\rho_n, t_n, u_n, \mathcal{H}_n) = ((\rho_{n-1})_1, (t_{n-1})_1, (u_{n-1})_1, (\mathcal{H}_{n-1})_1)$$

from the previous lemma and the discussion preceding it.

Let $\mathcal{H}' = \overline{\bigcup_{n \geq 1} \mathcal{H}_n}$ and define $\rho' = \varinjlim \rho_n$, $t' = \varinjlim t_n$ and $u' = \varinjlim u_n$. By [26, Theorem 3.3] $(\rho', t', \mathcal{H}')$ is an isometric representation of (X, \mathcal{C}) . Note that u' is a strongly continuous unitary representation of \mathcal{G} since it is the direct sum of such representations.

Now to the covariance relations:

$$\begin{aligned} u'(s)\rho'(c)P_{\mathcal{H}_n} &= u_n(s)\rho_n(c)P_{\mathcal{H}_n} \\ &= \rho_n(\alpha_s(c))u_n(s)P_{\mathcal{H}_n} \\ &= \rho'(\alpha_s(c))u'(s)P_{\mathcal{H}_n} \end{aligned}$$

for all $s \in \mathcal{G}, c \in \mathcal{C}$ and $n \in \mathbb{N}$. As well,

$$\begin{aligned} u'(s)t'(\xi)P_{\mathcal{H}_n} &= u_{n+1}t_{n+1}(\xi)P_{\mathcal{H}_n} \\ &= t_{n+1}(\alpha_s(\xi))u_{n+1}(s)P_{\mathcal{H}_n} \\ &= t_{n+1}(\alpha_s(\xi))u_n(s)P_{\mathcal{H}_n} \\ &= t'(\alpha_s(\xi))u'(s)P_{\mathcal{H}_n} \end{aligned}$$

for all $s \in \mathcal{G}, \xi \in X$ and $n \in \mathbb{N}$. Therefore, the covariance relations are satisfied and $(\rho', t', u', \mathcal{H}')$ is an isometric coextension of $(\rho, t, u, \mathcal{H})$.

Now let

$$\mathcal{K} = \overline{\text{span}}\{t'(\xi_1) \cdots t'(\xi_n)h : \xi_i \in X, h \in \mathcal{H}, n \geq 1\} \subset \mathcal{H}'.$$

Because of the covariance relations of $(\rho', t', u', \mathcal{H}')$ one can see that \mathcal{K} is a reducing subspace of $(\rho', t', u', \mathcal{H}')$ that contains \mathcal{H} . Thus, $(\rho', t', u', \mathcal{K})$ is a minimal isometric coextension of $(\rho, t, u, \mathcal{H})$.

In regard to uniqueness, suppose that $(\rho'', t'', u'', \mathcal{H}'')$ is another isometric coextension of $(\rho, t, u, \mathcal{H})$. [26, Proposition 3.2] proves that there exists a unitary $W : \mathcal{K} \rightarrow \mathcal{H}''$ such that $W\rho'(\cdot) = \rho''(\cdot)W$, $Wt'(\cdot) = t''(\cdot)W$ and $Wh = h$, for all $h \in \mathcal{H}$. Now

$$\begin{aligned} Wu'(s)t'(\xi_1) \cdots t'(\xi_n)h &= Wt'(\alpha_s(\xi_1)) \cdots t'(\alpha_s(\xi_n))u'(s)h \\ &= t''(\alpha_s(\xi_1)) \cdots t''(\alpha_s(\xi_n))Wu(s)h \\ &= t''(\alpha_s(\xi_1)) \cdots t''(\alpha_s(\xi_n))u(s)h \\ &= t''(\alpha_s(\xi_1)) \cdots t''(\alpha_s(\xi_n))u''(s)Wh \\ &= u''(s)t''(\xi_1) \cdots t''(\xi_n)Wh \\ &= u''(s)Wt'(\xi_1) \cdots t'(\xi_n)h \end{aligned}$$

Therefore, by minimality $Wu'(\cdot) = u''(\cdot)W$ and so $(\rho', t', u', \mathcal{K})$ and $(\rho'', t'', u'', \mathcal{H}'')$ are unitarily equivalent. \blacksquare

Theorem 4.7. *Let (X, \mathcal{C}) be a non-degenerate C^* -correspondence and let α be a generalized gauge action of a locally compact group \mathcal{G} . Then*

$$\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{T}_X^+ \rtimes_{\mathcal{T}_X, \alpha} \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha} \mathcal{G}}^+$$

Proof. It is already proven in [20], in a discussion following Theorem 7.13, that $\mathcal{T}_X^+ \rtimes_{\mathcal{T}_X, \alpha} \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha} \mathcal{G}}^+$. (It also follows from [3, Theorem 3.1] as the isomorphism Φ of that theorem maps generators to generators.)

Towards proving the remaining isomorphism, let $\varphi : \mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \rightarrow B(\mathcal{H})$ be a completely contractive representation. In the same way as in the proof of [20, Theorem 4.1] one can assume that φ is nondegenerate. Now by [20, Proposition 3.8] there exists a representation (π, u, \mathcal{H}) of $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ so that $\varphi = \pi \rtimes u$.

By [26, Theorem 3.10] there is a completely contractive representation, (ρ, t) , of (X, \mathcal{C}) such that $\pi = \rho \rtimes t$. Hence, in the same way as the first part of the proof of Theorem 4.4, $(\rho, t, u, \mathcal{H})$ is a completely contractive representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$. By Theorem 4.6 $(\rho, t, u, \mathcal{H})$ has a unique minimal isometric coextension $(\rho', t', u', \mathcal{H}')$ and thus by Theorem 4.4 $(\rho' \rtimes t', u', \mathcal{H}')$ is an isometric representation of $(\mathcal{T}_X, \mathcal{G}, \alpha)$.

As discussed in the proof of [26, Theorem 3.10] $\mathcal{H} \subset \mathcal{H}'$ is semi-invariant for ρ' and t' and thus $P_{\mathcal{H}}\rho' \rtimes t'|_{\mathcal{H}}$ is a completely contractive representation of \mathcal{T}_X^+ . Moreover, that same theorem gives that

$$\pi = \rho \rtimes t = P_{\mathcal{H}}\rho' \rtimes t'|_{\mathcal{H}}$$

because $\rho(c) = P_{\mathcal{H}}\rho'(c)|_{\mathcal{H}}$ and $t(\xi) = P_{\mathcal{H}}t'(\xi)|_{\mathcal{H}}$ for all $c \in \mathcal{C}$ and $\xi \in X$.

Therefore, every completely contractive representation of $\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G}$ dilates to a completely contractive representation of $\mathcal{T}_X^+ \rtimes_{\mathcal{T}_X, \alpha} \mathcal{G}$ and thus they are completely isometrically isomorphic. \blacksquare

Corollary 4.8. *Let $((X, \mathcal{A}), \mathcal{G}, \alpha)$ be a C^* -correspondence dynamical system and assume that $\mathcal{I}_X = \{0\}$. Then all relative crossed products for $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ are canonically isomorphic via completely isometric maps.*

In particular, the above applies to the non-commutative disc algebra $\mathcal{A}_{\infty} \subseteq \mathcal{O}_{\infty}$. To obtain the same conclusion for the non-commutative disc algebras \mathcal{A}_n , $n < \infty$, we need to work much harder. (See the next section.)

For the moment, we can put together all previous results to obtain the following, which summarizes our knowledge on the Hao-Ng isomorphism problem for the full crossed product.

Theorem 4.9. *Let $((X, \mathcal{A}), \mathcal{G}, \alpha)$ be a non-degenerate C^* -correspondence dynamical system. Then the following statements are successively weaker*

- (i) $C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G}) \simeq \mathcal{O}_X \rtimes_{\alpha} \mathcal{G}$ via a $*$ -isomorphism that sends generators to generators.
- (ii) $\mathcal{O}_X \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{O}_{X \rtimes_{\alpha} \mathcal{G}}$ via a $*$ -isomorphism that sends generators to generators. (Hao-Ng isomorphism)
- (iii) all relative crossed products for $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ are completely isometrically isomorphic via canonical maps.

If (X, \mathcal{C}) is hyperrigid, e.g., X is countably generated and $\varphi_X(\mathcal{J}_X)X = X$, then all of the above statements are equivalent.

Proof. Assume first that

$$(11) \quad C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G}) \simeq \mathcal{O}_X \rtimes_{\alpha} \mathcal{G}$$

canonically. Theorem 4.7 shows now that

$$\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{T}_X^+ \rtimes_{\mathcal{T}_X, \alpha} \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha} \mathcal{G}}^+$$

canonically and so by taking C^* -envelopes we have a canonical isomorphism

$$(12) \quad C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G}) \simeq C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\mathcal{T}_X, \alpha} \mathcal{G}) \simeq \mathcal{O}_{X \rtimes_{\alpha} \mathcal{G}}.$$

By “equating” the right sides of (11) and (12), we obtain (ii).

Assume now that (ii) is valid, i.e., the Hao-Ng isomorphism is implemented via a canonical map. The same map establishes

$$(13) \quad \mathcal{T}_X^+ \rtimes_{\mathcal{O}_X, \alpha} \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha} \mathcal{G}}^+.$$

By Theorem 4.7 we also have

$$(14) \quad \mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{T}_{X \rtimes_{\alpha} \mathcal{G}}^+.$$

From (13) and (14), we obtain $\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{T}_X^+ \rtimes_{\mathcal{O}_X, \alpha} \mathcal{G}$, or,

$$\mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G} \simeq \mathcal{T}_X^+ \rtimes_{C_{\text{env}}^*(\mathcal{T}_X^+), \alpha} \mathcal{G}$$

canonically. By Theorem 2.4, all relative crossed products for $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ are canonically isomorphic, which is (iii).

Assume now that (X, \mathcal{C}) is hyperrigid and all relative crossed products for $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ are canonically isomorphic. Therefore (7) in Theorem 3.6 implies

$$C_{\text{env}}^*(\mathcal{T}_X^+ \rtimes_{\mathcal{O}_X, \alpha} \mathcal{G}) \simeq \mathcal{O}_X \rtimes_{\alpha} \mathcal{G}.$$

By assumption $\mathcal{T}_X^+ \rtimes_{\mathcal{O}_X, \alpha} \mathcal{G} \simeq \mathcal{T}_X^+ \rtimes_{\alpha} \mathcal{G}$ and so (i) is valid.

Finally recall that Theorem 3.1 shows that a countably generated C^* -correspondence X with $\varphi_X(\mathcal{J}_X)X = X$ is always hyperrigid. \blacksquare

The importance of the previous result can not be understated. First, according to condition (iii), the verification of the Hao-Ng isomorphism for hyperrigid C^* -correspondences depends on the canonical identification of two non-selfadjoint operator algebras. We pursue this direction successfully in the next section where we verify the Hao-Ng isomorphism for all graph correspondences of row finite graphs. Furthermore, unlike condition (ii) (Hao-Ng isomorphism), both conditions (i) and (iii) are applicable to *arbitrary* dynamical systems $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$, i.e., α does not have to be a gauge action. Thus in a sense, a generalization of the Hao-Ng isomorphism problem beyond the realm of gauge actions is possible but only in the language of non-selfadjoint operator algebras. In light of the recent results of Harris and Kim [14], this seems to be a direction worth pursuing.

5. GRAPH CORRESPONDENCES

Following from the last section, we would like to prove that every isometric (Toeplitz) representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$ dilates to a covariant (Cuntz-Pimsner) representation. However, the standard proofs that the C^* -envelope of the tensor algebra is the Cuntz-Pimsner algebra are non-constructive [19, 26] which at the moment is a barrier to our method of proof. Significantly, in the case of graph correspondences such a constructive dilation proof is shown to exist.

Let (E, V, s, r) be a directed graph, where both E and V are separable, with associated graph correspondence $(X, \mathcal{C}, \varphi_X)$. Recall this is where $\mathcal{C} = c_0(V)$, X is the completion of $c_c(E)$ under the right module structure

$$\langle \alpha \delta_e, \beta \delta_f \rangle = \begin{cases} \bar{\alpha} \beta \delta_{s(e)}, & e = f \\ 0, & \text{otherwise} \end{cases}$$

$$\delta_e \cdot \delta_v = \begin{cases} \delta_e, & s(e) = v \\ 0, & \text{otherwise} \end{cases} ,$$

and the left action of \mathcal{C} on X is given by

$$\varphi_X(\delta_v) \delta_e = \begin{cases} \delta_e, & r(e) = v \\ 0, & \text{otherwise} \end{cases} .$$

When (X, \mathcal{C}) is the graph correspondence for a directed graph (V, E) then \mathcal{O}_X is $*$ -isomorphic to the Cuntz-Krieger algebra of the graph.

We wish to find Cuntz-Pimsner representations of these graph correspondences. As usual, the main concern is looking at which elements of \mathcal{C} are mapped into $\mathcal{K}(X)$ by φ_X .

In the case of a graph correspondence, Raeburn [29, Proposition 8.8] gives that $\varphi_X(\delta_v) \in \mathcal{K}(X)$ if and only if $|r^{-1}(v)| < \infty$. Furthermore,

$\delta_v \in \ker \varphi_X$ if and only if $r^{-1}(v) = \emptyset$. Thus, let

$$V_{\text{fin}} = \{v \in V : 1 \leq |r^{-1}(v)| < \infty\}$$

be the set of vertices generating Katsura's ideal \mathcal{J}_X and let

$$\mathcal{K} = \{(v, w) \mid \exists e \in E \text{ with } s(e) = w, r(e) = v, v \in V_{\text{fin}}\}.$$

For each pair $(v, w) \in \mathcal{K}$, let $E(v, w)$ be the collection of all edges starting from w and ending on v and let $[E((v, w))] = \mathbb{C}^{|E(v, w)|}$. In what follows we will identify the canonical basis of $[E(v, w)]$ with the elements of $E(v, w)$ and use the same symbol for both.

Suppose α is a generalized gauge action of (X, \mathcal{C}) then it is clear that this induces a permutation of V and in particular that $V_{\mathcal{K}}$ is invariant under this permutation. By abuse of notation we call this permutation $\alpha : V \rightarrow V$. Furthermore, the action α maps $[E(v, w)]$ unitarily onto $[E(\alpha(v), \alpha(w))]$. Indeed, if $E(v, w) = \{e_1, e_2, \dots, e_n\}$ and $\xi = \sum_{i=1}^n c_i \delta_{e_i}$, then

$$\begin{aligned} \langle \alpha(\xi), \alpha(\xi) \rangle &= \left\langle \alpha \left(\sum_{i=1}^n c_i \delta_{e_i} \right), \alpha \left(\sum_{j=1}^n d_j \delta_{e_j} \right) \right\rangle \\ &= \alpha \left(\left\langle \sum_{i=1}^n c_i \delta_{e_i}, \sum_{j=1}^n d_j \delta_{e_j} \right\rangle \right) \\ &= \alpha \left(\sum_{i=1}^n |c_i|^2 \delta_w \right) \\ &= \sum_{i=1}^n |c_i|^2 \delta_{\alpha(w)} = \langle \xi, \xi \rangle. \end{aligned}$$

Proposition 5.1. *Let (X, \mathcal{C}) be the graph correspondence of (E, V) and suppose $(\rho, t, u, \mathcal{H})$ is a completely contractive representation of the dynamical system $((X, \mathcal{C}), \mathcal{G}, \alpha)$. There exists a dilation to a completely contractive representation $(\rho_1, t_1, u_1, \mathcal{H}_1)$ such that for every $v \in V_{\text{fin}}$*

$$\rho(\delta_v) = \sum_{e \in r^{-1}(v)} t_1(\delta_e) t_1(\delta_e)^*$$

Proof. By [26, Lemma 3.5] because (ρ, t, \mathcal{H}) is completely contractive then for $v \in V_{\mathcal{K}}$ we have the matrix inequality

$$[t(e)^* t(f)]_{e, f \in r^{-1}(v)} \leq [\rho(\langle e, f \rangle)]_{e, f \in r^{-1}(v)} = \oplus_{e \in r^{-1}(v)} \rho(\delta_{s(e)})$$

and so $[t(e) : e \in r^{-1}(v)]$ is a row contraction. Hence,

$$\rho(\delta_v) \geq \sum_{e \in r^{-1}(v)} t(\delta_e) t(\delta_e)^*$$

and so we can define

$$\Delta_v := \frac{1}{\sqrt{|r^{-1}(v)|}} \left(\rho(\delta_v) - \sum_{e \in r^{-1}(v)} t(\delta_e) t(\delta_e)^* \right)^{1/2}.$$

Let $\mathcal{H}_w = \rho(\delta_w)\mathcal{H}$ and so we can assume $\mathcal{H} = \oplus_{w \in V} \mathcal{H}_w$. For each pair $(v', w') \in \mathcal{K}$ let

$$\mathcal{H}_{v', w'} := \mathcal{H}_{v'} \otimes [E(v', w')].$$

Then for each $w \in V$ we define

$$\mathcal{H}_w^+ \equiv \bigoplus_{(v', w) \in \mathcal{K}} \mathcal{H}_{v', w} = \bigoplus_{(v', w) \in \mathcal{K}} \mathcal{H}_{v'} \otimes [E(v', w)].$$

and

$$\mathcal{H}_{w, 1} = \mathcal{H}_w \oplus \mathcal{H}_w^+.$$

We combine these to define

$$\mathcal{H}_1 = \oplus_{w \in V} \mathcal{H}_{w, 1}.$$

Hence

$$\rho_1(\delta_w) = I_{\mathcal{H}_{w, 1}}, \quad w \in V,$$

extends to a *-homomorphism of \mathcal{C} that dilates ρ .

We also have a continuous unitary representation $u_1 : \mathcal{G} \rightarrow B(\mathcal{H}_1)$ dilating $u : \mathcal{G} \rightarrow B(\mathcal{H})$ and defined as follows.

Given $g \in \mathcal{G}$ and $h \in \mathcal{H}$, we let $u_1(g)h = u(g)h$. Otherwise, on each \mathcal{H}_w^+ the operator $u_1(g)$ is defined by

$$\mathcal{H}_w^+ \supseteq \mathcal{H}_{v', w} \ni h \otimes \xi \longmapsto \alpha_g(h) \otimes \alpha_g(\xi) \in \mathcal{H}_{\alpha_g(v'), \alpha_g(w)} \subseteq \mathcal{H}_{\alpha(w)}^+.$$

It is easy to see that $u_1 : \mathcal{G} \rightarrow \mathcal{B}(\mathcal{H}_1)$ is a continuous unitary representation dilating u .

We are ready to dilate $t : X \rightarrow B(\mathcal{H})$. If $r(e) \notin V_{\text{fin}}$, then we let $t_1(e) = t(e)$. Otherwise, if $e \in E$ with $s(e) = w$ and $r(e) = v \in V_{\text{fin}}$, then $t_1(e) \in B(\mathcal{H}_1)$ has cokernel contained in

$$\mathcal{H}_w \oplus (\dots 0 \oplus 0 \oplus \mathcal{H}_{v, w} \oplus 0 \dots) \subseteq \mathcal{H}_w \oplus \mathcal{H}_w^+ \equiv \mathcal{H}_{w, 1}$$

range contained in $\mathcal{H}_v \oplus 0 \subseteq \mathcal{H}_v \oplus \mathcal{H}_v^+$ and it is given by

$$t_1(\delta_e) = [t(\delta_e) \quad \Delta_v \tau(e)] \in B(\mathcal{H}_w \oplus \mathcal{H}_{v, w}, \mathcal{H}_v),$$

where

$$\tau(e) : \mathcal{H}_{v, w} \longrightarrow \mathcal{H}_v; \quad h \otimes \xi \longmapsto \langle e, \xi \rangle h.$$

(In general, for $\zeta \in [E(v, w)]$, $\tau(\zeta)$ will be given by $\tau(\zeta)(h \otimes \xi) = \langle \zeta, \xi \rangle h$.)

It is easy to see that $\tau(e)\tau(e)^* = I_{\mathcal{H}_v}$ and so,

$$\begin{aligned} \sum_{e \in r^{-1}(v)} t_1(\delta_e)t_1(\delta_e)^* &= \sum_{e \in r^{-1}(v)} t(\delta_e)t(\delta_e)^* + \Delta_v \tau(e)\tau(e)^* \Delta_v \\ &= \left(\sum_{e \in r^{-1}(v)} t(\delta_e)t(\delta_e)^* \right) + |r^{-1}(v)| \Delta_v^2 \\ &= \rho(\delta_v) \end{aligned}$$

This establishes one of the main conclusions of this proposition and gives that

$$[t_1(e)^*t_1(f)]_{e,f \in r^{-1}(v)} \leq [\rho_1(\langle e, f \rangle)]_{e,f \in r^{-1}(v)}$$

as in the start of the proof. Thus, by [26, Lemma 3.5] again, $(\rho_1, t_1, \mathcal{H}_1)$ is a completely contractive representation of (X, \mathcal{C}) .

Lastly we must establish the covariance relations. For any $g \in \mathcal{G}$ recall that α_g acts as a unitary between $[E(v, w)] = [\{e_1, e_2, \dots, e_n\}]$ and $[E(\alpha_g(v), \alpha_g(w))] = [\{f_1, f_2, \dots, f_n\}]$. This implies that

$$\begin{aligned} \sum_{i=1}^n t(\alpha_g(\delta_{e_i}))t(\alpha_g(\delta_{e_i}))^* &= \sum_{i=1}^n t\left(\sum_{j=1}^n (\alpha_g)_{j,i} \delta_{f_j}\right) t\left(\sum_{k=1}^n (\alpha_g)_{k,i} \delta_{f_k}\right)^* \\ &= \sum_{j,k=1}^n \left(\sum_{i=1}^n (\alpha_g)_{j,i} \overline{(\alpha_g)_{k,i}} \right) t(\delta_{f_j})t(\delta_{f_k})^* \\ &= \sum_{j=1}^n t(\delta_{f_j})t(\delta_{f_j})^* \end{aligned}$$

and so

$$\begin{aligned} u(g)(\rho(\delta_v) - \sum_{e \in r^{-1}(v)} t(\delta_e)t(\delta_e)^*) &= (\rho(\delta_{\alpha_g(v)}) - \sum_{e \in r^{-1}(v)} t(\alpha_g(\delta_e))t(\alpha_g(\delta_e))^*)u(g) \\ &= \left(\rho(\delta_{\alpha_g(v)}) - \sum_{w \in s(r^{-1}(v))} \sum_{e \in E(v,w)} t(\alpha_g(\delta_e))t(\alpha_g(\delta_e))^* \right) u(g) \\ &= \left(\rho(\delta_{\alpha_g(v)}) - \sum_{w \in s(r^{-1}(\alpha_g(v)))} \sum_{f \in E(\alpha_g(v),w)} t(\delta_f)t(\delta_f)^* \right) u(g) \\ &= \left(\rho(\delta_{\alpha_g(v)}) - \sum_{f \in r^{-1}(\alpha_g(v))} t(\delta_f)t(\delta_f)^* \right) u(g). \end{aligned}$$

By a standard functional analysis trick,

$$u(g)\Delta_v = \Delta_{\alpha_g(v)}u(g).$$

Furthermore, if $h \otimes \xi \in \mathcal{H}_{v,w}$, then

$$\begin{aligned} \tau(\alpha_g(e))u_1(g)(h \otimes \xi) &= \tau(\alpha_g(e))(u(g)h \otimes \alpha_g(\xi)) \\ &= \langle \alpha_g(e), \alpha_g(\xi) \rangle u(g)h \\ &= \langle e, \xi \rangle u(g)h = u(g)\tau(e)(h \otimes \xi). \end{aligned}$$

Hence,

$$\begin{aligned} t_1(\alpha_g(\delta_e))u_1(g) &= [t(\alpha_g(\delta_e)) \ \Delta_{\alpha_g(v)}\tau(\alpha_g(e))] (u_1(g) |_{\mathcal{H}_w \oplus \mathcal{H}_w^+}) \\ &= [u(g)t(\delta_e) \ \Delta_{\alpha_g(v)}u(g)\tau(e)] \\ &= [u(g)t(\delta_e) \ u(g)\Delta_v u(g)\tau(e)] \\ &= u_1(g)t_1(\delta_e). \end{aligned}$$

It is also immediate that

$$u_1(g)\rho_1(\delta_v) = \rho_1(\delta_{\alpha_g(v)})u_1(g).$$

Therefore, $(\rho_1, t_1, u_1, \mathcal{H}_1)$ is a completely contractive representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$. \blacksquare

Theorem 5.2. *Let (X, \mathcal{C}) be the graph correspondence of (E, V) . Every completely contractive representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$ can be dilated to a Cuntz-Pimsner representation.*

Proof. Let $(\rho_0, t_0, u_0, \mathcal{H}_0)$ be a completely contractive representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$. Recursively use the previous proposition to generate a sequence of completely contractive representations $(\rho_n, t_n, u_n, \mathcal{H}_n)$ such that for each $n \geq 1$, $\mathcal{H}_{n-1} \subset \mathcal{H}_n$, $(\rho_n, t_n, u_n, \mathcal{H}_n)$ is a dilation of $(\rho_{n-1}, t_{n-1}, u_{n-1}, \mathcal{H}_{n-1})$ and for every $v \in V$ such that $1 \leq |r^{-1}(v)| < \infty$ we have

$$\sum_{e \in r^{-1}(v)} t_n(\delta_e)t_n(\delta_e)^* = \rho_{n-1}(\delta_v).$$

Thus, define $\mathcal{H}' = \cup_{n=0}^{\infty} \mathcal{H}_n$ and

$$\rho'(c)|_{\mathcal{H}_n} = \rho_n(c), t'(\xi)|_{\mathcal{H}_n} = t_n(\xi), \text{ and } u'(g)|_{\mathcal{H}_n} = u_n(g).$$

Hence, $(\rho', t', u', \mathcal{H}')$ is a dilation $(\rho_0, t_0, u_0, \mathcal{H}_0)$ to a completely contractive representation such that for every $v \in V_{\text{fin}}$

$$\begin{aligned} \sum_{e \in r^{-1}(v)} t'(\delta_e)t'(\delta_e)^* &= \text{SOT} - \lim_{n \rightarrow \infty} \sum_{e \in r^{-1}(v)} t'(\delta_e)I_{\mathcal{H}_n}t'(\delta_e)^* \\ &= \text{SOT} - \lim_{n \rightarrow \infty} \sum_{e \in r^{-1}(v)} t_n(\delta_e)t_n(\delta_e)^* \\ &= \text{SOT} - \lim_{n \rightarrow \infty} \rho_{n-1}(\delta_v) \\ &= \rho'(\delta_v). \end{aligned}$$

According to [26, Definition 5.3] the representation $(\rho', t', \mathcal{H}')$ is \mathcal{J}_X -coisometric, i.e., Cuntz-Pimsner in the sense of Katsura but without being isometric.

Lastly, by Theorem 4.6 there is a unique minimal isometric coextension of $(\rho', t', u', \mathcal{H}')$ to $(\rho'', t'', u'', \mathcal{H}'')$. By [26, Corollary 5.21] this coextension process preserves the property of being \mathcal{J}_X -coisometric. Therefore, $(\rho'', t'', u'', \mathcal{H}'')$ is an isometric and \mathcal{J}_X -coisometric dilation of $(\rho_0, t_0, u_0, \mathcal{H}_0)$. We have therefore obtained a Cuntz-Pimsner representation of $((X, \mathcal{C}), \mathcal{G}, \alpha)$. ■

We now obtain the following which is valid for *arbitrary* graph correspondences.

Corollary 5.3. *Let (X, \mathcal{C}) be the graph correspondence of a directed graph (E, V) and $((X, \mathcal{C}), \mathcal{G}, \alpha)$ a C^* -correspondence dynamical system, that is, α is a generalized gauge action. Then all relative crossed products for $(\mathcal{T}_X^+, \mathcal{G}, \alpha)$ are canonically isomorphic via completely isometric maps.*

Proof. This is an immediate consequence of Theorem 5.2 and Theorem 2.4. ■

With an added assumption we obtain a positive solution to the Hao-Ng isomorphism problem.

Corollary 5.4. *Let (X, \mathcal{C}) be the graph correspondence of a row-finite directed graph (E, V) and $((X, \mathcal{C}), \mathcal{G}, \alpha)$ a C^* -correspondence dynamical system. Then*

$$\mathcal{O}_{X \rtimes_{\alpha} \mathcal{G}} \simeq \mathcal{O}_X \rtimes_{\alpha} \mathcal{G}.$$

Proof. By the description of \mathcal{J}_X mentioned at the start of this section and Theorem 3.1, the graph correspondence of a row-finite directed graph (E, V) is hyperrigid. The conclusion follows from Theorem 4.9. ■

Acknowledgement. Part of this research was carried out in the summer of 2018 while EK was visiting Chongqing Normal University, Chongqing University, the Chinese Academy of Science in Beijing and the Chern Institute of Mathematics. EK would like to thank his hosts Professors Liming Ge, Hanfeng Li, Chi-Keung Ng, Wenhua Qian, Wenming Wu and Wei Yuan for the stimulating conversations and their hospitality during his stay at their institutions.

REFERENCES

- [1] B. Abadie, *Takai duality for crossed products by Hilbert C^* -bimodules*, J. Operator Theory **64** (2010), 19–34.
- [2] W. Arveson, *The noncommutative Choquet boundary II: hyperrigidity*, Israel J. Math. **184** (2011), 349–385.
- [3] E. Bedos, S. Kaliszewski, J. Quigg and D. Robertson, *A new look at crossed product correspondences and associated C^* -algebras*, J. Math. Anal. Appl. **426** (2015), 1080–1098.
- [4] E. Bedos, S. Kaliszewski, J. Quigg and J. Spielberg, *On finitely aligned left cancellative small categories, Zappa-Szj products and Exel-Pardo algebras*, manuscript, arXiv:1712.09432.
- [5] D. Blecher and C. Le Merdy, *Operator algebras and their modules-an operator space approach*, London Mathematical Society Monographs New Series **30**, Oxford University Press, 2004.
- [6] N. Brown and N. Ozawa, *C^* -algebras and finite-dimensional approximations*, Graduate Studies in Mathematics **88**, American Mathematical Society, Providence, RI, 2008. xvi+509 pp.
- [7] V. Deaconu, *Group actions on graphs and C^* -correspondences*, Houston J. Math. **44** (2018), 147–168.
- [8] V. Deaconu, A. Kumjian and J. Quigg, *Group actions on topological graphs*, Ergodic Theory Dynam. Systems **32** (2012), 1527–1566.
- [9] B. Duncan, *Certain free products of graph operator algebras*, J. Math. Anal. Appl. **364** (2010), 534–543.
- [10] R. Exel, *Amenability for Fell bundles*, J. Reine Angew. Math. **492** (1997), 41–73.
- [11] N. Fowler, *Discrete product systems of Hilbert bimodules*, Pacific J. Math. **204** (2002), 335–375.
- [12] E. Gootman and A. Lazar, *Crossed products of type I AF C^* -algebras by abelian groups*, Isr. J. Math. **56** (1986), 267–279.
- [13] G. Hao and C-K. Ng, *Crossed products of C^* -correspondences by amenable group actions*, J. Math. Anal. Appl. **345** (2008), 702–707.
- [14] S. Harris and S. Kim, *Crossed products of operator systems*, manuscript arXiv:1803.10759.
- [15] E.T.A. Kakariadis, *The Dirichlet property for tensor algebras*, Bull. Lond. Math. Soc. **45** (2013), 1119–1130.
- [16] S. Kaliszewski, J. Quigg and D. Robertson, *Functoriality of Cuntz-Pimsner correspondence maps*, J. Math. Anal. Appl. **405** (2013), 1–11.
- [17] S. Kaliszewski, J. Quigg and D. Robertson, *Coactions on Cuntz-Pimsner algebras*, Math. Scand. **116** (2015), 222–249.
- [18] E. Katsoulis, *C^* -envelopes and the Hao-Ng isomorphism for discrete groups*, Inter. Math. Res. Not. (2017), 5751–5768.
- [19] E. Katsoulis and D. Kribs, *Tensor algebras of C^* -correspondences and their C^* -envelopes*, J. Funct. Anal. **234** (2006), 226–233.
- [20] E. Katsoulis and C. Ramsey, *Crossed products of operator algebras*, Memoirs Amer. Math. Soc, to appear.
- [21] E. Katsoulis and C. Ramsey, *Crossed products of operator algebras: applications of Takai duality*, J. Funct. Anal. (2018), to appear.

- [22] D.W. Kim, *Coactions of Hopf C^* -algebras on Cuntz-Pimsner algebras*, arXiv:1407.6106 [math.OA].
- [23] D. Larson and B. Solel, *Structured triangular limit algebras* Proc. London Math. Soc. **75** (1997), 177–193.
- [24] M. McAsey and P. Muhly, *Representations of nonselfadjoint crossed products*, Proc. London Math. Soc. **47** (1983), 128–144.
- [25] A. Morgan, *Cuntz-Pimsner algebras and twisted tensor products*, arXiv:1601.07826.
- [26] P. Muhly and B. Solel, *Tensor algebras over C^* -correspondences: representations, dilations, and C^* -envelopes*, J. Funct. Anal. **158** (1998), 389–457.
- [27] V. Paulsen, *Completely Bounded Maps and Operator Algebras*, Cambridge Studies in Advanced Mathematics **78**, Cambridge University Press, 2002. MR1976867
- [28] G. Popescu, *Isometric dilations for infinite sequences on noncommuting operators*, Trans. Amer. Math. Soc. **316** (1989), 523–536.
- [29] I. Raeburn, *Graph algebras*, CBMS Regional Conference Series in Mathematics **103**, American Mathematical Society, Providence, RI, 2005.
- [30] J. Schaëffer, *On unitary dilations of contractions*, Proc. Amer. Math. Soc. **127** (1955), 322.
- [31] C. Schafhauser, *Cuntz-Pimsner algebras, crossed products and K -theory*, J. Funct. Anal. **269** (2015), 2927–2946.
- [32] D. Williams, *Crossed products of C^* -algebras*, Mathematical Surveys and Monographs, Vol. 134, American Mathematical Society, 2007.

DEPARTMENT OF MATHEMATICS, EAST CAROLINA UNIVERSITY, GREENVILLE,
NC 27858, USA

E-mail address: katsoulise@ecu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, MACEWAN UNIVERSITY,
EDMONTON, AB, CANADA

E-mail address: ramseyc5@macewan.ca